

**SECURING A FUTURE FOR
FISH AND WILDLIFE**

A Conservation Legacy for Iowans

**Author:
James J. Zohrer
E Resources Group**

**Edited by:
Dr. Terry W. Little
Iowa DNR
Wildlife Research Supervisor**

**Douglas C. Harr
Iowa DNR
Wildlife Diversity Program Coordinator**

**Jeffrey R. Vonk, Director
Iowa Department of Natural Resources
September 2005
Revised May 2006**

Table of Contents

	<u>Page</u>
Table of Contents	i
List of Tables	iv
Lists of Figures	v
List of Maps	v
List of Appendices	vii
Letter from the Director	viii
Foreword from the Steering Committee	ix
Acknowledgements	xi

Securing a Future for Fish and Wildlife - *A Conservation Legacy for Iowans*

Chapter One: Introduction, Purpose and Planning	1
Introduction	1
Purpose	3
The Planning Process	4

THE STATUS OF IOWA'S FISH AND WILDLIFE IN 2005

Chapter Two: The State of Iowa's Natural Communities	11
Physiography	11
Historic Plant Communities	15
Historic Wildlife Communities	18
Impacts of Settlement	20
Change Continues in the 20 th Century	24
Wildlife Conservation	30
Iowa's Natural Communities Today	32

Chapter Three: Species of Greatest Conservation Need	34
Wildlife Species considered by the IWAP	34
Determining the Species of Greatest conservation Need	34
SGCN in Each Taxonomic Class	51
Chapter Four: Habitats of SGCN	52
Terrestrial Habitat Classes	52
Aquatic Habitat Classes	65
Habitat Preferences of SGCN	69
Priorities for Habitat Protection	71
Chapter Five: Stresses on Iowa's Wildlife and Its Habitats	73
Direct Stresses on Wildlife	73
Stresses on Wildlife Habitats	74
Stress Analysis	78
Stresses on Terrestrial Wildlife and Habitats	79
Stresses on Aquatic Wildlife and Habitats	92
<i>The Status of Iowa's Fish and Wildlife in 2005 -</i>	98
 Summary and Conclusion	

THE IOWA WILDLIFE ACTION PLAN

Chapter Six: A Vision for Iowa in the Year 2030	101
A Vision for Iowa's Wildlife	103
A Vision for Wildlife Habitats	104
A Vision for Wildlife Management	106
A Vision for Wildlife-Associated recreation	108
A Vision for Wildlife Education	109
A Vision to Fund Wildlife Education	111
Chapter Seven: Research, Surveys and Monitoring	112
General Discussion	112
New Survey Needs	114
Long Term Monitoring	114
Objective 1: Current Inventory of Wildlife in Iowa	118
Objective 2: Monitoring Species and Their Habitats	120
Objective 3: Strengthening Vertebrate GAP Models	123
Objective 4: Impact Assessment	123

Chapter Seven: (Continued)	
Objective 5: Evaluation of Management Protocols and Restoration Programs	124
Research Priorities	125
Research and Monitoring costs	126
Chapter Eight: Priorities for Conservation Actions	137
General Discussion	137
Priorities for Vision Elements	129
Wildlife Vision	129
Habitat Vision	134
Management Vision	147
Recreation vision	147
Education Vision	149
Funding vision	150
Chapter Nine: Plan Coordination, Implementation and Review	151
Plan Coordination	151
Plan Implementation	151
Interagency Cooperation	153
IWAP Review	155
Chapter Ten - The Cost of Sustaining Iowa's Biodiversity	156
Appendices 1 - 21	159

List of Tables

Chapter One: Introduction, Purpose and Planning

Table 1-1. IWAP Steering Committee	5
Table 1-2. IWAP Steering Committee Working Groups	6
Table 1-3. Iowa Steering Committee Meetings	7

Chapter Three: Species of Greatest Conservation Need

Table 3-1. Number of species considered by the IWAP	34
Table 3-2. Breeding birds of greatest conservation need	38
Table 3-3. Migratory birds of greatest conservation need	40
Table 3-4. Mammals of greatest conservation need	41
Table 3-5. Reptiles and amphibians of greatest conservation need	42
Table 3-6. Land snails of greatest conservation need	43
Table 3-7. Butterflies of greatest conservation need	44
Table 3-8. Fish of greatest conservation need	46
Table 3-9. Mussels of greatest conservation need	48
Table 3-10. Dragonflies and damselflies of greatest conservation need	50
Table 3-11. Percent of Iowa species in each taxonomic class listed as a SGCN	51

Chapter Four: Habitats of SGCN

Table 4-1. Description of Terrestrial Habitat Classes Used in the IWAP	52
Table 4-2. GIS Land Cover Types From 2002 Satellite Imagery	53
Table 4-3. The Amount of Iowa's Wildlife Habitat Found in Each Landform	54
Table 4-4. Landcover Within Each Landform	54
Table 4-4. Aquatic Habitat Classes Used in the IWAP	65
Table 4-5. Lake types by landform	68
Table 4-6. Summary of Habitat Preferences of SGCN by Habitat Class	70

Chapter Five: Stresses on Iowa's Wildlife and Its Habitats

Table 5-1. Definitions of Stress Levels	78
Table 5-2. Terrestrial stresses affecting Iowa's wildlife and its habitats	80
Table 5-3. Aquatic stresses affecting Iowa's wildlife and its habitats	81
Table 5-4. Stresses on Terrestrial Habitats (averaged over all taxonomic groups and all landforms)	82
Table 5-5. Stresses on Terrestrial Wildlife (averaged over all habitat classes)	85
Table 5-6. Statewide Stresses on Birds	86
Table 5-7. Statewide Stresses on Mammals	87

Table 5-8. Statewide Stresses on Reptiles and Amphibians	88
Table 5-9. Stresses on Land Snails (found only in Paleozoic Plateau)	89
Table 5-10. Statewide Stresses on Butterflies	90
Table 5-11. Statewide Stresses on Fish and Mussels	93
Table 5-12. Statewide Stresses on Dragonflies and Damselflies	97
Chapter Seven: Research, Surveys and Monitoring	
Table 7-1. Summary of Wildlife Monitoring, Surveys and Inventories Conducted in Iowa: 1960-2005	113
Table 7-2. Short Term Survey Needs for SGCN	115
Table 7-3. Land Management Research Needs	121
Table 7-4. Wildlife Species Research Needs	122
Table 7-5. Area-Specific Research Projects	123
Table 7-6. Estimated research and monitoring costs	124
Chapter Eight: Priorities for Conservation Actions	
Table 8-1. Priority Habitats by Landform	146
Chapter Ten - The Cost of Sustaining Iowa's Biodiversity	
Table 6-1. Cost to Double the Amount of Permanently Protected Conservation Land in Iowa by 2030	157

List of Figures

Chapter One: Introduction, Purpose and Planning	
Figure 1-1. Sales of Iowa Hunting Licenses	2
Chapter Two: The State Of Iowa's Natural Communities	
Figure 2-1. Changes In Cropping Patterns and Representative Grassland Wildlife	26
Figure 2-2. Statewide Pheasant Trends	27
Figure 2-3. Trends in Bobwhite Quail and Harvest	29

List of Maps

Chapter Two: The State of Iowa's Natural Communities	
Map 2-1. Landform Regions of Iowa	12
Map 2-2. Landcover of Iowa in the 1850's	16
Map 2-2. Publicly-owned Conservation Lands in Iowa	31
Map 2-3. Landcover of Iowa in 2002	33

List of Maps (continued)

Chapter Four: Habitats of SGCN	
Map 4-1. Forest Land Cover	56
Map 4-2. Wet Forest/Woodland Land Cover	57
Map 4-3. Woodland Land Cover	58
Map 4-4. Shrubland Land Cover	59
Map 4-5. Wet Shrubland Land Cover	60
Map 4-6. Herbaceous Wetland Land Cover	61
Map 4-7. Warm Season Herbaceous Land Cover	62
Map 4-8. Savanna Land Cover	63
Map 4-9. Agricultural Land Cover.	64
Map 4-10. Major Lakes and River Systems of Iowa	67
Chapter Five: Stresses on Iowa's Wildlife and Its Habitats	
Map 5-1. Predominant Land Use by Watershed	94
Map 5-1. Iowa's Impaired Waters	95
Chapter Eight: Priorities for Conservation Actions	125
Map 8-1. Restorable Wetland Soils in the Prairie Pothole Region of Iowa	128
Map 8-2. All Terrestrial Vertebrate Species Richness	130
Map 8-3. Bird Species Richness	130
Map 8-4. Mammal Species Richness	131
Map 8-5. Reptile Species Richness	131
Map 8-6. Amphibian Species Richness	132
Map 8-7. Aquatic Species Richness	132
Map 8-8. Existing Protected Land Complexes of 2,000 Acres or larger	135
Map 8-9. Prairie Pothole Joint Venture Priority Wetland Complexes	137
Map 8-10. Landowner Incentive Program Site Priorities	138
Map 8-11. The Nature Conservancy's Priority Conservation Areas	139
Map 8-12. Existing and Proposed Bird Conservation Areas	140
Map 8-13. Iowa Audubon's Important Bird Areas	141
Map 8-14. Topeka Shiner Critical Habitat	142
Map 8-15. Migration Habitat Across Northern Iowa	143
Map 8-16. WRP Special Project Areas	144
Map 8-17. High Priority Areas for Cooperative Conservation Actions	145
Map 8-18. Distribution of Iowa's Human Population	148
Map 8-19. Distribution of Existing Public Lands	149

List of Appendices

Appendix 1.	The IWAP - A Plan To Plan	159
Appendix 2.	State Advisory Group Meeting Summary	164
Appendix 3.	List Of Iowa Breeding Birds Considered By The IWAP.	182
Appendix 4.	List Of Migratory Birds Considered By The IWAP.	189
Appendix 5.	List Of Iowa Mammals Considered By The IWAP.	194
Appendix 6.	List Of Iowa Fish Considered By The IWAP.	197
Appendix 7.	List Of Iowa Amphibians And Reptiles Considered By The IWAP	201
Appendix 8.	List Of Iowa Mussels Considered By The IWAP.	204
Appendix 9.	List Of Iowa Butterflies Considered By The IWAP.	206
Appendix 10.	List Of Iowa Dragonflies And Damselflies Considered By The IWAP.	210
Appendix 11.	NatureServe National And Subnational (State) Heritage Status Rank Definitions.	213
Appendix 12.	Iowa Animal Endangered, Threatened, Or Species Of Special Concern	215
Appendix 13.	Terrestrial Habitat Classes Used In The IWAP.	219
Appendix 14.	Habitat Preferences For Species Covered By The IWAP	226
Appendix 15.	SGCN Grouped By Habitat Class	244
Appendix 16.	Regional Stress Analysis Tables Terrestrial Habitats	266
Appendix 17.	Aquatic Habitat Stresses By Region	291
Appendix 18.	Iowa Lakes And Rivers By Region	301
Appendix 19.	Existing Large Habitat Complexes In Public Ownership By Region	310
Appendix 20.	References Cited And Used In Document Preparation	319
Appendix 21.	Monitoring And Surveys Conducted On Iowa's Wildlife: 1960-2004	326



September 9, 2005

Dear Iowans:

This document - *The Iowa Wildlife Action Plan* - is a guide for the conservation of Iowa's wildlife for the next 25 years. Two years in the making, it is the product of a cooperative effort between more than 100 professional fish and wildlife biologists, educators, land managers and concerned citizens who provided the guidance and information essential to its development.

This is Iowa's first truly comprehensive plan designed to aid all wildlife - from birds to freshwater mussels and fish to butterflies. Fully one-third of Iowa's native fish and wildlife is declining rapidly as a result of habitat loss and declining water quality. This plan will improve the status of these declining species and at the same time benefit common wildlife that we all know and enjoy.

This is a habitat-oriented plan that when completed will double the amount of permanently protected wildlife habitat in the state. It will help preserve Iowa's precious wildlife heritage, provide recreation benefits to our citizens, improve our quality of life and increase the economic activity from bird watching, hiking, nature photography, fishing, hunting and many other wildlife-associated activities..

When you read the plan I hope you will share my excitement for the potential it expresses for fish and wildlife and the citizens that enjoy them. Please join me in enthusiastically endorsing this plan and support the funding necessary to see that it becomes a reality.

Sincerely,

Jeffrey R. Vonk
Director

Foreword from the Steering Committee

In October 2003, the Iowa Department of Natural Resources (IDNR) assembled the nucleus of a steering committee to develop a Comprehensive Wildlife Conservation Plan for Iowa. The thought of creating such a plan was daunting. We were charged with developing a plan that would surpass anything ever prepared for the conservation of Iowa's wildlife. A comprehensive review of the status and needs of *all* of Iowa's wildlife had never been attempted. The Iowa Board of Conservation, drawing on the work of Aldo Leopold, created *The Iowa Twenty-five Year Conservation Plan* for game species in 1933. That plan did not begin to address the status of, stresses to, or needs of all the great diversity of vertebrate and invertebrate wildlife. Game species management plans have been used by IDNR for years, but the only other wildlife plans developed in Iowa were restoration plans for selected individual species or for rare and endangered habitats.

We committee members represent the IDNR, non-governmental conservation organizations, academia, and private enterprise. All have some special expertise and a special interest in the future of Iowa's wildlife. We recognized from the start that this plan must be useful not just to IDNR, but to every individual and organization concerned about wildlife in Iowa.

To help us with this monumental task we added members and assembled working teams of specialists to deal with various faunal groups and other pertinent issues. Work teams coordinated with the Plan Author, Plan Coordinator, and Steering Committee Chair to provide species lists and environmental stresses, review draft chapters, and assist with a public Advisory Group meeting held in July, 2004.

We want this document to become more than just another plan to put on a shelf and ignore. It is our greatest wish that it will become the *Iowa Wildlife Action Plan*—a living document that can be updated and used to help guide wildlife management in Iowa for many years. The steering committee hopes you may find this prospect as exciting as we do.

Douglas C. Harr
Steering Committee Chair

Terry W. Little
Plan Coordinator

James J. Zohrer
Plan Author

The Steering Committee:

Todd Bishop

Kim Bogenschutz

Don Brazelton

Angi Bruce

Dave DeGeus

James J. Dinsmore

Bruce Ehresman

Marlene Ehresman

Dale Garner

Barb Gigar

Daryl Howell

Karen Kinhead

Erwin Klaas

Monica Ulman

Terry VanDeWalle

Michelle Wilson

Ric Zarwell

Guy Zenner

Acknowledgements

This Plan could not have been completed without the assistance of dozens of individuals that provided advice, data, coordination, writing skills and review of the Plan in its many stages. Special recognition is due the members of the Plan Steering Committee without whose leadership, guidance and hard work completion of the Plan would not have been possible: Don Brazelton (Iowa Association of County Conservation Boards), David DeGeus (The Nature Conservancy), Marlene Ehresman (Iowa Natural Heritage Foundation), Dr. Ervin Klaas and Dr. James Dinsmore (Iowa State University, retired). Terry VanDeWalle (EarthTech, Inc.), Ric Zarwell (Iowa Audubon), and Department of Natural Resource employees Richard Bishop, Dr. Terry Little, Dr. Dale Garner, Douglas Harr, Todd Bishop, Kim Bogenschutz, Bruce Ehresman, Angi Bruce, Daryl Howell, and Monica Ulman.

In addition, 93 individuals representing 59 government or private conservation organizations, educational institutions, agricultural support groups and private wildlife-associated enterprises served on the Plan Advisory Committee. The Advisory Committee developed the Plan's vision and strategies and provided comments on draft versions.

Twenty-six other individuals served on 13 Working Groups that did the extensive resource reviews and discussions necessary to determine the Species of Greatest Conservation Need.

U.S. Fish and Wildlife Service employees Michael Sweet and Dr. John Christian provided helpful comments on an earlier version of the Plan. Mike Sweet interpreted Federal planning guidelines, assisted with administrative issues and provided contact and resource information that made the entire planning process easier.

Thirty individuals provided comments on the first complete review draft of the Plan. Dr. James Dinsmore (Iowa State University, retired), Jeff Lerner (Defenders of Wildlife), Jim Munson (U.S. Fish and Wildlife Service) and Jeff Kopaska (IDNR Fisheries Bureau) provided especially thoughtful reviews.

DNR employee Kerri Wells designed the cover.

Dr. Terry Little and Doug Harr edited and compiled the final document.

James Zohrer
Plan Author
E Resources Group

Chapter One

Introduction, Purpose and Process


Introduction

Fish and wildlife conservation programs in Iowa have been funded nearly entirely from license fees paid by hunters and anglers and by Federal excise taxes collected on hunting and fishing equipment. Iowa Department of Natural Resources (IDNR) conservation efforts have historically featured game animals in spite of the fact that just 147 of the 999 species addressed in this plan are hunted, fished or trapped. Wildlife professionals have long recognized that nongame wildlife (animals that are not fished, hunted or trapped) is an important component of natural ecosystems and that their status is often an indicator of general environmental health. IDNR has been involved for 25 years in the restoration of charismatic nongame like river otters, Trumpeter swans, peregrine falcons and others (See Chapter 2) in an attempt to generate a broader funding base for nongame programs. Without a dedicated funding source, however, most nongame species have benefited only from game-oriented programs that have coincidentally restored or improved habitat for a wide range of species.

The number of hunting licenses sold in Iowa has declined 45 percent over the last 60 years (Figure 1-1) and fishing license sales are decreasing at 1 percent annually. This decline in field sports has been blamed on the rural-to-urban migration of the past century and to competition for citizen's time in our increasingly busy, technology-driven lives. Whatever the reason, this decline in license buyers foretells a potential drop in financial support for wildlife programs if the trends continue. Increases in license fees and the growth of deer and turkey hunting have increased IDNR revenues and masked this erosion in underlying support. Evidence suggests this growth has stabilized and is likely to decrease as baby boomers reach retirement, and probable retirement from hunting and angling. The funding available to maintain wildlife programs into the future could be very much in jeopardy unless alternate sources of funding are found.

Significant national efforts to find funding for nongame wildlife programs began during the last quarter of the 20th century. Congress passed the Nongame Act in 1980 to appropriate funds to the states for nongame management. In 1981 the Iowa legislature initiated a state income tax check-off (Chickadee Checkoff) to provide matching funds for anticipated Federal grants. The Nongame Act was never funded, however, and check-off funds have never been adequate to address the many needs of nongame species. Currently the Chickadee Check-off generates approximately \$150,000 annually, compared to \$30 million generated by hunters and anglers.

Figure 1-1. Sales of Iowa hunting licenses



In 1989 the International Association of Fish and Wildlife Agencies began a nationwide campaign to develop stable, sufficient, long-term funding for nongame programs. *Teaming With Wildlife* rallied state and territorial fish and wildlife agencies to develop grassroots coalitions of nongame enthusiasts that would lobby Congress for funds. The proposed funding mechanism was a Federal excise tax on nongame-related materials and equipment that would have been apportioned to the states in the same manner as the taxes collected on hunting and angling equipment. This effort was opposed by the outdoor recreation industry and failed to gain support in Congress.

The *Conservation and Reinvestment Act (CARA)* was proposed in 1996 as a second attempt to find funding for nongame. Offshore oil and gas lease funds would be permanently redirected from the Federal treasury to state accounts that would have paid for a variety of wildlife, cultural and environmental protection programs. CARA would have generated \$350 million nationwide annually for wildlife; \$4.5 million would have been Iowa's share. CARA passed in the House of Representatives and had widespread support and momentum in the Senate when the terrorist activities of September 11, 2001, redirected Federal budget priorities. Permanent funding failed to pass, but the *Wildlife Conservation and Restoration Program (WCRP)* was enacted as a one-time appropriation for state nongame programs. Similar acts titled *State and Tribal Wildlife Grants (SWG)* have been passed annually since then. These grants

have required non-federal matching funds that vary from 25% to 50% depending on the year and type of program.

Purpose

As a condition of receiving SWG funds, Congress has mandated that state fish and wildlife agencies develop a *Comprehensive Wildlife Conservation Plan* (Plan) by October 1, 2005. The Plan must include eight required elements:

1. Information on the distribution and abundance of wildlife that IDNR deems to be indicative of the diversity and health of Iowa's wildlife. Low and declining populations of fish and wildlife are defined in the Plan as Species of Greatest Conservation Need (SGCN)
2. Locations and relative conditions of key habitats and community types essential to conservation of SGCN.
3. Problems that adversely affect SGCN or their habitats and priority research and survey efforts needed to identify factors that may assist in their restoration and improved conservation.
4. Conservation actions necessary to conserve SGCN and their habitats and priorities for implementing such actions.
5. Plans for monitoring SGCN and their habitats, for monitoring the effectiveness of conservation actions, and for adapting these conservation actions to new information or changing conditions.
6. Procedures to review the Plan at intervals not to exceed ten years.
7. Plans for coordinating, to the extent feasible, the development, implementation, review, and revision of conservation actions with Federal, state, and local agencies that manage significant land and water areas in Iowa, or administer programs that significantly affect the conservation of SGCN and their habitats.
8. Descriptions of public participation in the development, revision, and implementation of the plan.

The Plan must utilize the best available knowledge on the distribution and abundance of wildlife, historical documentation and other references to identify Iowa's wildlife conservation needs. The Plan must address the needs of all wildlife, but focus primarily on SGCN and their habitats as determined by IDNR.

The Planning Process

Steering Committee. Responsibility for developing the IWAP was vested by Congress in the Iowa IDNR, and by the IDNR in the department's Wildlife Bureau. To be sure that the Plan had the widest possible acceptance, the bureau identified a Steering Committee composed of employees and volunteers from government and nongovernmental conservation organizations (NGOs), educational institutions and the private sector (Table 1-1). Steering Committee members had demonstrated expertise in the many disciplines necessary for developing a Plan of this magnitude.

The Steering Committee developed a planning schedule with milestones, selected a plan author and assigned responsibilities for accomplishing specific tasks (Appendix 1). The first step was to describe the distribution and abundance of wildlife, identify SGCN and their habitats, and identify problems affecting them. Thirteen Working Groups led by steering committee members performed this time-consuming work (Table 1-2). These Plan elements are contained in the chapters 2 - 5 of this Plan titled *Iowa's Fish and Wildlife and Their Habitats - History and Status In 2005*

To provide a focus for the planning efforts, the steering committee made the following decisions:

- The IWAP would be a wildlife plan. Plants are not specifically discussed except in habitat-related sections.
- The IWAP would have a 25-year focus. Long-term continuity was needed to accomplish ambitious objectives, but achievements needed to be accomplished in a time frame that could be appreciated by Plan supporters.
- The IWAP would be strategic in nature. Operational plans to implement its vision and strategies would be crafted later to fit the unique missions and capabilities of conservation organizations and individuals interested in implementing them.

Much of the work undertaken by the Steering Committee and Working Groups was conducted through e-mail, regular mail and the telephone. Steering Committee meetings were held to coordinate Plan development, team updates, status reports, reviews and other activities (Table 1-3).

Table 1-1. IWAP Steering Committee

Name	Title	Affiliation	Plan Responsibilities
Richard Bishop	Wildlife Bureau Chief Retired 9/04	IDNR	Plan Director
Dr. Dale Garner	Wildlife Bureau Chief Effective 9/04	IDNR	Plan Director
Dr. Terry Little	Wildlife Research and Wildlife Diversity Supervisor	IDNR	Plan Coordinator
Doug Harr	Wildlife Diversity Program Manager	IDNR	Steering Committee Chair
Todd Bishop	GIS Systems Supervisor	IDNR	GIS data
Don Brazelton	Executive Secretary	Iowa Association of County Conservation Boards	County Conservation Boards - funding - legislation
Kim Bogenschutz	Fisheries Biologist	IDNR	Aquatics (fish and invertebrates)
David DeGeus	Director of Conservation Programs	The Nature Conservancy	TNC input and NGO's
Bruce Ehresman	Wildlife Diversity Biologist	IDNR	Birds, wildlife diversity programs
Marlene Ehresman	Program/Planning Associate	Iowa Natural Heritage Foundation	Landscapes and habitats
Dale Garner (Replaced by Guy Zenner) 11/04	Executive Officer Waterfowl Biologist	IDNR	Federal plan coordination
Angi Bruce	Wildlife District Supervisor	IDNR	Wildlife management and private lands
Daryl Howell	Senior Environmental Specialist	IDNR	T & E species and small mammals
Dr. Ervin Klaas	Professor Emeritus	Iowa State University	GAP, mammals, terrestrial invertebrates
Terry VanDeWalle	Biologist	EarthTech, Inc.	Reptiles and Amphibians
Ric Zarwell	Important Bird Area Coordinator	Audubon	Birds
Monica Ulman	GIS Specialist	IDNR	GIS data and mapping

Table 1-2. IWAP Steering Committee Working Groups

Working Group Leaders	Responsibilities	Others Having Initial Input
Dr. Terry Little	Game Species	Todd Bogenschutz - IDNR Peter Fritzell - IDNR
Daryl Howell and Dr. Ervin Klaas	Mammals	Dr. John Bowles, Professor Emeritus, Central College
Kim Bogenschutz	Fish and Mussels	Daryl Howell John Olson - IDNR, Environmental Specialist Tom Wilton - IDNR, Environmental Specialist Kelly Poole - IDOT, Aquatic Ecologist Scott Gritters - IDNR, Fisheries Biologist
Bruce Ehresman	Birds	Dr. James Dinsmore, Professor Emeritus, ISU Ric Zarwell Doug Harr Robert Cecil - Author, Iowa Ornithologists Union (IOU) Ann Johnson - IOU Darwin Koenig - Author Bob Russell - U.S. Fish and Wildlife Service
Terry VanDeWalle	Reptiles and Amphibians	Dr. J.L. Christiansen -Professor, Drake University
Daryl Howell	Butterflies	Jerry Selby - Private Consultant Dr. Diane Debinski - Professor, ISU
Daryl Howell	Land Snails	
Dr. Erv Klaas	Dragonflies and Damselflies	Ann Johnson - IOU
Dr. Erv Klaas	GAP	Kevin Kane - ISU
Todd Bishop	GIS	Monica Ulman
Barb Gigar Michelle Wilson	Education and Recreation	IDNR Fisheries Education IDNR Planning and Coordination
Dr. Karen Kinkead	Monitoring and Evaluation	Dr. Erv Klaas, Bruce Ehresman, Kim Bogenschutz, David DeGeus
Angi Bruce	Management and Habitat	Brent Olson -IDNR Forestry Scott Moats - TNC Kevin Pape - IDNR Parks Monica Ulman - IDNR - Wildlife Kelly Smith - IDNR - Wildlife Dan Cohen - Buchanan CCB Tom Gengerke - IDNR - Fisheries Joe Boyles - Polk CCB Marlene Ehresman

Table 1-3. Iowa Steering Committee Meetings

2003	2004	2005
July 1, 2003	February 5, 2004	January 18, 2005
October 23, 2003	April 27, 2004	April 6, 2005
December 3, 2003	August 10, 2004	
	September 21, 2004	
	November 9, 2004	

Advisory Group. To assure the Plan would involve a diversity of conservation viewpoints, representatives of 105 conservation, recreation, education and agricultural support organizations were invited to serve on a formal Advisory Group; 93 individuals representing 59 organizations agreed to participate (Appendix 2-1).

The Advisory Group met in Des Moines on July 17, 2004. The purpose of the meeting was to develop a vision for the IWAP and strategies for attaining that vision by the year 2030. The Advisory Group was updated on the planning process and the status of wildlife and their habitats in Iowa (Appendices 2-2 and 2-3). The large group then broke into eight focus groups and developed vision elements and conservation actions (Appendices 2-3 and 2-4). When condensed by the steering committee, these vision elements and conservation actions form the basis for the second part of this Plan - the *Iowa Conservation Action Plan* (IWAP)(Chapters 6 - 10).

Stresses. The steering committee and fisheries and wildlife research and management biologists from the IDNR identified stresses affecting Iowa's wildlife in each of 8 major landform regions. Terrestrial and aquatic stresses were evaluated and ranked for each of the major taxonomic classes included in the Plan (Chapter 5 and Appendices 16 and 17).

Other planning. To complete the Plan, new Working Groups (Table 1-2) were formed by the steering committee to develop priorities for research and monitoring needs, identify priorities for conservation actions and site specific opportunities for Plan implementation, and develop education and recreation strategies.

One of the key factors identified during the process of determining the SGCN was the lack of current, credible information on the distribution and abundance of many nongame species. Iowa GAP provides habitat-based distribution information, but much of the information used in GAP models is old or anecdotal - little is based on formal wildlife surveys. A new Working Group identified research and monitoring needs and proposed a model for statewide evaluation of wildlife populations as the Plan is implemented (Chapter 7).

A Working Group utilized information from Iowa GAP and DNR GIS coverage maps to identify regions of the state with the greatest potential for protection of key habitats (Chapter 8). These maps also indicate regions where habitat is lacking and restoration programs are needed to improve wildlife populations.

Two additional required Plan elements are priorities for implementing conservation actions and plans for coordination and review. These are covered in Chapters 8 and 9, respectively.

Estimated costs for implementing the IWAP are presented in Chapter 10.

Public participation. Persons with much of the ecological and conservation expertise existing in the state were included in various stages of the planning process, either as formal participants or consultants and reviewers on specific portions of the IWAP. A variety of efforts were made to ensure that information about the Plan received statewide distribution to the public as well:

- A popular article - *Iowa Wildlife Species of Greatest Conservation Need* - was published in the January/February 2005 issue of the *Iowa Conservationist* magazine. The *Conservationist* has a circulation of 25,000 including every school and public library in the state, and a readership of 100,000 individuals.
- A section on the IDNR's Wildlife Diversity web site provided an overview of the IWAP.
- From March 2004 through June 2005, 20 formal presentations were given on the Plan that reached approximately 1,500 individuals. These included IDNR staff, affiliated conservation agencies, NGOs and professional wildlife groups throughout the state.

The first complete draft of the IWAP was finished in April 2005, and was made available to the public in several ways:

- A complete draft and executive summary were placed on the IDNR's web site with email links to a post office that received comments.
- Paper copies or a CD were supplied to readers upon request.
- Notice was sent directly to the 165 individuals representing 92 different organizations that were invited to the Advisory Group meeting (Appendix 2).
- Statewide news releases advertised completion of the Plan, where it was available and how to comment.

- The public comment period for the draft IWAP was held from May 1st through June 10th, 2005. A total of 30 written comments were received and incorporated in whole or part into the final version of the Plan.

While a large number of individuals contributed in some manner to the IWAP, ultimate responsibility for its content lies with the Steering Committee and the Iowa Department of Natural Resources.

IOWA'S FISH AND WILDLIFE
AND THEIR HABITATS
HISTORY AND STATUS IN 2005

CHAPTER TWO

THE STATE OF IOWA'S NATURAL COMMUNITIES

Physiography

Topography. Iowa is a state of 56,239 square miles bordered by the Mississippi River on the east, and the Missouri and Big Sioux Rivers on the west. Iowa has a relatively low relief - elevations run from a high of 1,670 feet above mean sea level in Osceola County in northwestern Iowa to 480 feet above mean sea level in Lee County in the southeastern corner of the state.

Climate. Iowa's climate is classified as humid continental and is characterized by warm summers and cold winters. A small portion of northern and northeastern Iowa are in the cool - summer subtype (Hillaker 1993). The average temperature in the summer ranges from 71 degrees F in the north to 73 degrees in the south part of the state. December to February winter temperatures average 22 degrees with an average winter difference of 6.5 degrees between north and south. Temperature minimums of -25 degrees F are not uncommon in northern Iowa.

Statewide winter snowfall averages 32 inches. Northern Iowa (north of U.S. Highway 30) receives frequent snow often associated with strong winds, blowing and drifting. Southern Iowa may experience substantial snowfall as well as more frequent ice storms. This results in a snow cover that is often covered by a surface crust of ice or hard snow. Harsh conditions seldom last for more than a few weeks in most of the state, even less in the south half.


The average annual precipitation is 32 inches. The northwest part of the state is the driest with an annual precipitation of 28 inches while the southeast is the wettest with an annual precipitation of 36 inches. Iowa often experiences seasonal extremes and frequent local, rapid weather changes due to the convergence of cold, dry Arctic air, moist maritime air from the Gulf of Mexico, and dry Pacific air masses. Like most states, periods of severe drought and periods of excessive precipitation can have a dramatic impact on terrestrial and aquatic vegetation as well as their associated fish and wildlife species.

These climatic factors combine to influence the length of the growing season across the state. Late frosts in the spring and early freezes in the fall result in a reduced growing season of 135 days in northeastern and northwestern Iowa. The longest growing season is in southeastern Iowa, with an average of 175 days. The state average growing season is 158 days long.

Geology. Iowa's natural communities are as much a result of its recent geologic past as they are a result of climatic conditions (Prior 1991). Prior (1991)

divides the state into seven geologic regions based on the underlying bedrock as well as the location of glacial and loess deposits. (Map 2-1). The boundaries of these landform regions coincide well with the boundaries of other habitat based classification systems. The italicized names for regions are those listed in Prior (1991). The names in parentheses are habitat-based names for the landforms that describe the native vegetation that was present at the time of settlement. Descriptions are taken from Prior (1991) and Iowa GAP.

Map 2-1. Landform Regions of Iowa



Source: Iowa DNR

The Loess Hills (Tallgrass Prairie) is a unique landform that formed at the end of the last Ice Age about 18,000 years ago. The formation is only one to fifteen miles wide but is about 200 miles long extending from near Sioux City, Iowa to St. Joseph, Missouri. Although deposits of windblown soils (loess) are found in many parts of the world, nowhere else but in China do they reach as high as in Iowa where some of the hills are more than 200 feet above the adjacent Missouri River valley. The Loess Hills landform has other features that are easily noticed. Bedrock is exposed naturally in only a few places and the soil

has unique physical properties. If the topsoil on the slope of a hill is removed, the exposed loess will erode quickly and deep gullies will form. Even when covered with topsoil, loess can slump, often in a unified way across a slope creating “cat-step” ledges along the sides of hills. However, when a loess hill is cut vertically the exposed wall will stand for decades.

The Des Moines Lobe (Prairie Potholes) has a landscape that is gently rolling with abundant moraines, shallow wetland basins or potholes, and a few relatively deep natural lakes. This landform still retains the imprints of recent glacial occupation. Loess is entirely absent. The most prominent landform patterns left by the Wisconsin glacier on the Des Moines Lobe are the end moraines. The Des Moines Lobe is part of the Prairie Pothole Region that extends north and west into western Minnesota, eastern North and South Dakota, and the Canadian Prairie Provinces. Most of the potholes have been drained with ditching and underground tile lines to make way for agriculture. Agriculture was also responsible for greatly increasing the rate at which streams and drainage patterns developed in this geologically young landform.

The **Southern Iowa Drift Plain (Tallgrass Prairie)** is the largest of Iowa’s landforms. Like the Des Moines Lobe, it is composed almost entirely of glacial drift, but the Pre-Illinoian glaciers that deposited material in this part of Iowa were much older. As a result, deep glacial drift, ranging from a few to several hundred meters, is the only evidence of their occupation. Instead of poorly drained and relatively level landscapes, streams have had time to erode the land surface and form well-defined drainage systems. Hilltops have similar elevations that reveal the approximate level of the land surface constructed by the last ice sheet. As erosion slowly dissected this landscape, a layer of loess ranging from 2 to 10 meters was deposited over the glacial till. Throughout the Southern Iowa Drift Plain the terrain varies considerably, but the pattern of relief resulting from its history of erosion is the dominant feature of the region. Many of the larger rivers had glaciers standing in their headwaters at the time the Des Moines Lobe was ice-covered. These valleys obtained much of their present width, depth, and alluvial fill from flooding as the Wisconsin ice sheet melted away from northcentral Iowa. In many places the rivers have cut through the glacial drift into the underlying sedimentary bedrock. The rough wooded terrain adjoining these valleys supports many scenic and recreational areas and important wildlife habitat.

The **Iowan Surface (Eastern Tallgrass Prairie)** landform extends over a large region of northeastern Iowa and is characterized by long, gently rolling slopes, low relief, and open views of the horizon. Pre-settlement vegetation in this region was primarily prairie with heavily wooded floodplains along the larger watercourses. Drainage networks are well developed, but stream gradients are low with some scattered areas of poor drainage and natural wetlands. The area was once part of the Pre-Illinoian Southern Iowa Drift Plain but experienced

large-scale and more destructive erosion events, the latest occurring during the coldest part of the Wisconsin glaciation 16,500 to 21,000 years ago. Frost action, down slope movement of water-soaked soil materials, and strong winds were the dominant geologic processes in this region. Layers of loess are thin and scattered. Glacial boulders are numerous and many are very large. Elongated ridges and isolated oblong hills called pahas occur in the southern part of the Iowa Surface region. These features are covered with a mantle of silt and sand believed to have accumulated in response to strong northwesterly winds that occurred during the period of glacial cold. Soils mapped on the larger pahas indicate they developed under forest vegetation rather than prairie. Karst topography occurs in the northern part of the landform where cavities in the underlying limestone bedrock collapsed and formed sinkholes. Fens are also present but more scattered than in the Des Moines Lobe.

The **Northwest Iowa Plain** (*Eastern Tallgrass Prairie*) contains many of the terrain features and geologic materials present in other landforms and is similar in appearance to the Iowa Surface with a uniform low relief. This landform was and still is a relatively treeless, gently rolling landscape. Despite these similarities, the landscape differs from other regions because of a combination of factors. The western uplands of this region are underlain with highly eroded, Pre-Illinoian glacial tills. The eastern part of these tills is covered with later glacial deposits from an early Wisconsin glacial advance. The entire region was then subjected to vigorous erosion activity that accompanied the later advance of the Wisconsin ice sheet. As a result, features of a freshly glaciated landscape were lost as a well-established branching network of streams formed over the entire region. The deeper thickness of the loess mantle, the overall elevation of the land surface, and the present precipitation and vegetation distinguish the Northwest Iowa Plains from the state's other landforms. Windblown loess is abundant and nearly continuous across the region ranging in thickness from 4 to 16 feet. Altitudes throughout the Northwest Iowa Plains are uniformly higher than any other portion of the state and topographically are continuous with the High Plains of the Dakotas. Average annual precipitation is lower than other parts of the state. Thus, the region is higher, drier and less timbered than any other in the state. Although bedrock exposures are rare in the Northwest Iowa Plains, the oldest bedrock in Iowa (Precambrian-age Sioux Quartzite) occurs here along the Big Sioux River.

The **Paleozoic Plateau** (*Prairie to Hardwood Transition*) is the most distinctive of Iowa's landforms because of its abundant rock outcroppings, karst topography, a near absence of glacial deposits, many deep narrow valleys, cool-water streams, and heavily wooded uplands. Numerous gorges and ravines cause abrupt local changes in the direction of slopes and exposures. These sites provide abundant cool, moist and wooded habitats rich in diverse communities of plants and animals. Seeps and springs are common features along valley sides where strata of varying permeability are exposed and signify subterranean drainage systems. Ice caves and cold-air (*algific*) slopes are unique to this area. Unusual

microclimates associated with these features support a particularly rare and sensitive biological habitat in Iowa. The steep rocky slopes are unsuited for agriculture and remain heavily forested. Remnant prairies occur on south and west facing slopes. Ecologists believe these prairies were more extensive before the suppression of naturally occurring fires following European settlement.

Alluvial Plains, often called floodplains, are constructed by water flowing off of the landscape and carrying with it boulders, cobbles, gravel, sand, silt, and clay. This process of erosion creates a dendritic-shaped landform of nearly level corridors with varying widths depending on the size and reach of the river. These corridors are largest along the Missouri and Mississippi Rivers but can be found along streams throughout the other landforms. The floodplain is a dynamic landform that is frequently disturbed, sometimes drastically, by flood and drought events. Stream channels may be cut off leaving backwater sloughs or oxbow lakes. Large-scale vertical changes may also occur within the floodplain due to the deposition of alluvium that forms terraces and benches. These structures are level but are elevated above existing floodplains by a distinct slope. Smaller tributaries that enter the floodplain of a larger river often form alluvial fans that may cover older floodplain materials. During low flow periods, wind becomes an important factor in the transport of materials. Exposed sand or soil having little or no vegetation to hold it place can be blown onto floodplain and terraces as well as onto higher elevations along valley margins. Sand dune topography occurs downwind of valley floors.


The glacial history and topography of each landform affect the type and distribution of current wildlife habitats and agricultural land use. These are discussed in Chapter 4.

Historic Plant Communities

Pre-settlement Iowa lay at a biological crossroads. Hardwood forests dominated the cooler and more humid lands east of the Mississippi River. The warmer, drier shortgrass prairie and prairie potholes of the northern Great Plains lay to the west. To the north, great maple-basswood and pine forests covered the Great Lakes region. To the south, oak savannas gradually gave way to the vast oak-hickory forests of the Missouri Ozarks. These different ecological regions blended together in Iowa to produce a unique landscape of great biological diversity (Map 2-2).

Roughly two-thirds of the state (an estimated 23 million acres) was dominated by lush prairies. Most was tallgrass prairie, although short grasses were present on hot, dry sites. Nearly 7 million acres of forest or forest-prairie savanna covered much of the eastern third of Iowa and followed the river valleys into the prairies to the north and west. Around 4 million acres of prairie

Map 2-2. Landcover of Iowa in the 1850's (from Government Land Office original public land survey of Iowa)



pothole marshes dotted recently-glaciated and poorly-drained northcentral and northwest Iowa. Another million acres of backwaters, sloughs and flooded oxbows were found in the floodplains of the Mississippi, Missouri and larger inland rivers.

Prairies. The prairie was more than just a monolithic sea of grass. Prairie plants are adapted to subtle changes in moisture and soils that occur along a gradient from lowlands to drier prairie ridges. Poorly drained wetlands and wetland margins supported rank growths of sedges, slough grass, cord grass, bluejoint, prairie muhly grass, and panic grass, with common forbs such as gayfeather, prairie dock, Turk's-cap lily and New England aster. Better-drained loamy soils on slopes and broad ridges were covered with more moderate stands of switchgrass, big bluestem, Indian grass and forbs like compass plant, rattlesnake master, smooth aster, wild indigo and goldenrod. Drier sites on gravel and sand ridges or steep slopes supported shorter and more open stands of little bluestem, side-oats grama, and needlegrass, with forbs like pasque flower, silky aster, yellow puccoon and common milkweed.

Forests. Closed-canopy mature forests as we know them today existed only on the floodplains where fire could not routinely penetrate. Silver maple, American elm, and swamp white oak dominated on the wettest sites, with hickories, hackberry, black walnut, white ash, red oak, basswood and slippery elm on lower slopes. Shrubs were not abundant and were primarily young silver maples and hackberry with catbriar, poison ivy and grape.

Forests on drier slopes and uplands were primarily oak openings or savannas - scattered old oak trees or small clumps of oaks with an understory of prairie or mixed prairie-forest shrubs and herbs. Burr oak, with its thick, fire-retardant bark dominated with some red and white oaks on moister sites. The understory was primarily prairie grasses and forbs but hazel, coralberry, sumac and grape occurred where fire was less common.

The heaviest concentrations of timber were in the cooler and moister eastern third of the state. In the west only the floodplains and the coolest sites on north and east facing slopes in the deepest river valleys were timbered. Because of the many river systems that penetrated the prairies to the north and west, however, at least some timber and shrub lands were found across most of the state.

Fire and grazing. Drought, fire and grazing combined to make Iowa's prairie-wetland-forest ecosystem an even more diverse place. In wet years water levels were high, wetland vegetation gradually died out, and marshes began to look like ponds or small lakes. But dry weather runs in approximately 10-year cycles on the prairies, with severe drought at roughly 20-year intervals. Drought caused wetland basins to temporarily de-water. Seeds buried in moist wetland soils were able to germinate once the water was removed and dense stands of

emergent vegetation were replenished. Thus regenerated wetlands awaited only the end of drought to return them to their former productive condition.

In wet years fire was less prevalent on the prairie. Without burning the dead stems and leaves of grasses and forbs accumulated on the ground and this litter created a cooler, moister environment. In some cases coralberry and other shrubs were able to survive and spread from forest edges farther into the grasslands. During drought fire burned off large areas of prairie and forest, killed invading shrubs and trees, eliminated the litter, returned nutrients to the soil and allowed grasses to regain their dominance. Fire also allowed annual plants like ragweed, fleabane, thistle and primrose to take a temporary foothold before the longer-lived grasses and forbs recovered and choked them out.

Although fires were common, it is impossible to say how much and how frequently the prairies burned. Weather is seldom in complete synchrony over all of Iowa. Local dry spells undoubtedly created mini-droughts that lowered wetlands and produced frequent fires, while just a few miles away precipitation was normal. Even in normal years a dry late summer could result in a partial drawdown of marshes and occasional fires. The network of wetlands, creeks and rivers probably stopped smaller fires from expanding too greatly.

Grazers and browsers like bison, elk and deer also had an affect. They suppressed shrubs and slowed the growth of tall grasses where they fed intensively. Elk and bison created wallows - sandy areas where they rolled in the loose earth to remove hair and discourage insects. Prairie dogs, though not common in Iowa, kept the vegetation around their towns clipped short. Even gophers created small openings over their mounds where annual plants could gain a foothold.

The result of all this variety in soils, topography, weather, fire and animal activity was a great patchwork of plant communities in both time and space. On some sites 250 species of plants could be found. Not only were prairies, forest and wetlands in close proximity, but at any given location plant communities were in a state of growth, retrenchment or suppression depending on their local history.

Historic Wildlife Communities

Game animals. The great diversity of plant communities that covered pre-settlement Iowa also supported a diversity and abundance of wildlife that was foreign to settlers from the East. Iowa native Aldo Leopold, writing in 1931 in his *Game Survey of the North Central States*, said, "...no region in the world was originally more richly endowed with game than this one, quantity and quality both considered. Contrary to common belief, the cream of its game country was the prairie type..." Prairie animals like bison, elk, pronghorn, prairie chickens and

sharp-tailed grouse penetrated the tallgrass prairies from the West. White-tailed deer, wild turkeys, passenger pigeons, bobwhite quail and ruffed grouse followed the deciduous woodlands and river valleys into the prairie from the East.

Waterbirds. The prairie pothole and riverine wetlands provided excellent nesting habitat and attractive resting and feeding stops for millions of migrating ducks and geese between their nesting and wintering grounds. Giant Canada geese, trumpeter swans and a dozen species of ducks nested in Iowa, mainly blue-winged teal, mallards, redheads, northern shovelers, northern pintails and ruddy ducks. In excess of 4 million ducks may have been raised annually. One early hunter thought there were 10,000 Canada goose nests in Kossuth County alone.

Other waterbirds were also plentiful. Sandhill cranes were abundant during migration and nested here occasionally. Whooping cranes were less numerous, but nested frequently in the marshes of northcentral and northwest Iowa. More than 30 species of shorebirds migrated through Iowa. Of these, the woodcock, long-billed curlew, marbled godwit and upland sandpiper nested here, and the golden plover, Eskimo curlew and common snipe were abundant during migration.

Furbearers. Beaver, muskrat and river otters were found throughout Iowa, associated entirely with marshes, streams and rivers. Muskrat were most abundant in the prairie marshes of northcentral Iowa and maintained very high numbers. Beaver and river otters were associated more with riparian habitats. Mink and raccoon were not highly sought after, but both must have been abundant.

Predators. An variety of predators fed on this abundance of game animals. The gray wolf occurred in two forms, the lighter, smaller Great Plains wolf followed the bison and elk herds and was most common in the western two-thirds of the state. The timber wolf, a somewhat larger and darker version of the same species, inhabited the forested eastern third, mostly in the northeast corner, and would have fed mainly on deer. Coyotes were found statewide. Red foxes were found in the prairies and at the prairie-forest border in northern Iowa, the gray fox was found primarily in the eastern third of the state. Bobcats were numerous, occurring statewide in a variety of forested and shrubby habitats. Mountain lions, or cougar, were occasionally seen, but reports are few and far between. The lynx, a larger version of the bobcat normally inhabiting the coniferous forests of the Great Lakes states and Canada, was at least occasionally found here.

The black bear was the largest predator in pre-settlement Iowa. Although their preferred habitat was woodlands, they occasionally wandered into the prairies. Reports of black bears originate from 48 counties, most in eastern Iowa.

The grizzly bear, arguably North America's most fierce predator, was found occasionally on the Great Plains, but there are no known records from Iowa.

Nongame. Records of the un hunted fauna that inhabited Iowa are largely nonexistent. The early explorers and settlers were concerned mostly with wildlife as a source of food or pelts, or as a threat to their livestock and crops. But of 440 species of birds and mammals that resided here or migrated through Iowa, less than 15 percent were ever hunted or trapped. Serious scientific efforts to describe Iowa's wildlife did not begin until nearly 40 years after settlement, and by then significant changes had already occurred.

The wetlands must have been home for yellow-headed blackbirds, marsh wrens, American bitterns, black terns, and Sora rails as well as Canada geese, mallards, and muskrats. Wetland-prairie margins must have been nesting sites for song sparrows, sedge wrens and northern harriers.

Where shrubby, early successional stages of forest pushed into the prairies there must have been an abundance of cardinals, yellowthroats, rufous-sided towhees and rose-breasted grosbeaks as well as ruffed grouse and white-tailed deer. Larger stands of mature forest must have provided nesting sites for ovenbirds, scarlet tanagers, wood thrushes and pileated woodpeckers as well as wild turkeys. Riparian woodlands would have been home for black-billed cuckoos, redheaded woodpeckers, belted kingfishers and northern flickers as well as wood ducks.

Grasshopper and vesper sparrows would have nested in recently burned prairies. Prairies a year or two after burning would have provided nesting cover for bobolinks and dickcissels as well as prairie chickens. Henslow's sparrows, savanna sparrows and upland sandpipers would have nested in older prairies with dense ground litter. Loggerhead shrikes and mourning doves would have sought out grasslands with a shrub component. In all, more than 180 species of birds nested in Iowa.

Even less is known of the reptiles, amphibians and invertebrates. Pristine prairie potholes, riverine wetlands, prairies and woodlands provided homes for a diversity of lizards like the Great Plains skink and six-lined racerunner, common turtles like the ornate, snapping and painted turtles, snakes like the timber and Mississauga rattlesnakes and frogs like the leopard, green and gray tree frog. More than 60 species of reptiles and amphibians were eventually found in Iowa.

Impacts of Settlement

Settlement in Iowa progressed roughly southeast to northwest. Most of the south half of the state had been inhabited by the end of the 1840's; northcentral and northwest Iowa were settled in the 1850's; Lyon County in

extreme northwest Iowa was the last to be settled, receiving its first homestead family in 1866.

Human population growth was slow at first. By 1840 only 43,000 settlers had braved the prairies. Pressure for cheap land increased after the Civil War, however, and massive land grants were made to railroad builders to stimulate completion of a transcontinental railroad network. By 1870, Iowa's population had skyrocketed to nearly 650,000, by 1900 to 2 million.

An Agricultural Revolution. At the same time Iowa was being settled a revolution was overhauling industry and agriculture:

- James Watt perfected the steam engine in 1790;
- Robert Fulton developed the first steam locomotive in the U. S. in 1802;
- The first steamboat plied the Mississippi River in 1811;
- A steam-powered dragline was used to finish the Erie Canal in 1825;
- Cyrus McCormick perfected a mechanical reaper in 1832;
- John Deere developed the first steel moldboard plow in 1836;
- By 1851 Deere was producing 10,000 steel plows a year and the fate of the prairie was sealed.
- In 1865 the first transcontinental railroad was completed and routes to the West were opened during the next 25 years.

The advent of improved farm implements, coupled with a rapidly expanding population base devoted mostly to agriculture, had a devastating and permanent impact on Iowa's native plant communities.

Forests. Woodlands were the first to go. Some early settlers preferred farming woodlands rather than open prairie, fearing that land too poor to grow trees would not grow crops either. While experience would quickly prove that wrong, forests felt the bite of the pioneer's ax early in our history.

Most of the initial forest clearing in Iowa was done to allow conversion of the land to agriculture. Iowa's native hardwoods did not prove valuable as building materials. Most of the lumber that eventually built the farm homes, barns and livestock dwellings that dotted the countryside came from the great pineries of Minnesota and Wisconsin. Starting in the 1850's, however, railroad expansion and the discovery of coal in southern Iowa fueled a demand for oak ties and mine timbers that would last into the early 20th century. By 1875, just one-third of the original 6.7 million acres of primitive forest remained, most on rough land or in floodplains either too steep or too wet to plow.

Prairies. The effect on our extensive prairies and prairie-wetland complexes was even more devastating. When pulled by up to 5 teams of horses or yokes of oxen a steel *breaking plow* could shear through and break up 2 acres a day of the foot-thick sod with its intricately intertwined root systems. The newly exposed soil was so fertile that a crop, first wheat and later corn, was planted

directly on the overturned furrows. The next year a second plowing would complete the conversion of prairie to a field tillable by conventional methods. Starting in the 1850's, Iowa lost nearly 2 percent of its 25 million acres of native prairie a year, 3 million acres a decade, until less than 30,000 acres (0.1%) remained after 80 years.

Wetlands. The vast prairie-pothole wetlands of northcentral and northwest Iowa took longer to impact. Through the first 20 years of settlement there was plenty of good land available without trying to farm around wet acres. In 1850, Congress passed the SwampLand Act. It directed each county to survey all wetlands and sell them at auction for 5 cents an acre, the first of what would become a century-long succession of government-subsidized efforts to drain wetlands. County drainage commissions and drainage districts were soon organized. Eventually pothole soils were discovered to be some of the most productive when dry, further accelerating the demand for drainage.

The first drainage attempts were with hand-dug, open ditches that drained shallow lakes and large marshes into permanent streams. Steam dredges did not replace hand labor until nearly 1900. Underground ceramic tiles were developed to drain smaller potholes into ditches as early as 1858. By 1917 modern clay tiles were used to drain seasonally wet fields into extensive, interconnected drainage systems that had eliminated all but the largest wetlands. By 1906 just 25 percent of the original 4 million acres of wetlands remained.

Wildlife. Iowa's original wildlife populations suffered a similar fate as her native plant communities. Species that competed with humans for space, or were particularly useful for food or fiber, or required very specific habitats that were eliminated or drastically reduced did not survive. Others of less importance to humans held on in low numbers wherever suitable habitat remained, but their future remained suspect. A few species that could adapt to the remarkable change in habitats or were favored by the conversion to a diversified agricultural environment increased for a time, at least until agriculture became too pervasive.

By 1900 the large game animals and the predators that lived on them were gone (bison, elk, white-tailed deer, gray wolves, mountain lions, black bear and bobcats). Smaller predators like coyotes and red and gray fox were more adaptable, fed on a wider range of smaller prey animals, and were able to survive in Iowa into the 20th century. Economically important furbearers like river otter and beaver were also essentially gone by 1900.

Wild turkeys, passenger pigeons, prairie chickens and waterfowl all fed occasionally on settler's crops and were considered pests, and all were valuable as table fare or to sell at local markets. The spread of railroads into the Midwest in the 1860's and 1870's allowed hunters to reach the best hunting grounds and permitted shipping frozen game to markets in Chicago, Milwaukee and as far as New York City before it spoiled. Game was served as a delicacy in many

eastern restaurants in the late 19th century. As city dwellers developed more leisure time in the 1880's, hunting for sport or recreation also became more popular.

The take of game birds was enormous. A single net could capture 1,500 passenger pigeons. Entire flocks of turkeys could be pot shot from the roost on cold winter nights. Hunters could occasionally take 100 or more prairie chickens in a day (seasonal takes of 900 or more chickens were recorded). Sport hunters were able to take up to 100 ducks in a single day. The best market hunters could take up to 3,0000 ducks in a season. One group of 7 hunters shipped 14,000 ducks east in a single year. A careful hunter willing to pick his shots could take a half dozen mallards or 8 or 9 prairie chickens with a single shot. Avid woodcock hunters could take 40 birds a day; one market hunter took up to 3,000 woodcock a year in northeast Iowa. A hunter could easily take several ruffed grouse in a day but apparently few were ever sold at market. A variety of shorebirds - long-billed and Eskimo curlews, marbled godwits, upland and golden plovers and common snipe - were frequently hunted and at least some sold at market. Whooping and Sandhill cranes were also hunted for the table and because they were a pest in grain fields.

For most of Iowa's early history this activity was totally unregulated. Seasons, bag limits, shooting hours and restrictions on weapons did not exist. Settlers shot game for the table year around as they could find it. Sport and market hunters were active primarily in Fall and Spring so that game would not spoil before it could be consumed or sold. By the 1880's market hunters were building freezers to prolong their hunting opportunities. Waterfowl suffered the additional indignity of having their eggs collected for food or by egg collectors, a common hobby in the later 1800's. There seemed to be no need for regulation - the game was limitless, far more than anyone could possibly use.

But as hunting pressure increased in the 1870's and 1880's, habitat loss was also accelerating. Iowa was becoming settled. Nearly every square mile of land had several farm families living on it. New farmers looked to more ways to create tillable land. Much of the forested land that remained into the 1870's was turned into pasture. Cattle, sheep and hogs destroyed the undergrowth and competed with wildlife for acorns and other native food. A variety of species that so far had been able to withstand the hunting pressure alone began to be affected by the increasing fragmentation and elimination of their habitats. Whatever the reason - unregulated hunting, habitat loss, or more likely a combination of both - much of the wildlife that had existed here for centuries was in severe decline by the late 1870's.

Ever smaller flights of passenger pigeons continued into the mid 1870's, dwindled more into the 1880's and 90's and were gone by 1900. Wild turkeys were gone from northeast Iowa by 1854, from most of central Iowa by the 1870's, and disappeared from southern Iowa by 1910. Ruffed grouse were able to hold

on into the 20th century only in the most heavily forested counties of northeast Iowa.

Prairie chickens and bobwhite quail fared somewhat better. Opening the prairies to grain farming provided an alternate winter food supply in grain stubble. More reliable foods allowed their numbers to increase and their range to expand as long as there was enough prairie remaining for nesting and winter cover. Prairie chicken numbers may have peaked in the 1870's. After that chickens and quail began declining as too much prairie was converted to crop fields. Both hung on at lower numbers well into the 20th century.

Waterfowl and shorebirds continued to migrate in large numbers through Iowa until the end of the 19th century. Fewer were produced here as prairies were turned over and wetlands drained, but spectacular migrations from breeding grounds on the prairies to the north undoubtedly softened the blow of local habitat loss. By the 1890's, however, the loss of wetlands was taking a toll and by 1900 market hunting was a thing of the past. The last Sandhill and Whooping crane nests were found in Hancock County in 1894, the last long-billed curlew nest in 1890, and the last giant Canada goose nest in 1910.

Clearing of forests, conversion of native prairies to farm fields and the draining of wetlands eliminated far more species of songbirds, reptiles and amphibians than just the game species. Most of the loss went unnoticed by settlers, and by the time the first naturalists began studying the flora and fauna of Iowa, much change had already occurred and went unrecorded.

In less than a century the landscape of Iowa was changed more by settlement than that of any other state. Most of Iowa's 2 million residents lived on small, nearly self-sufficient farms of 100 acres or less. They subsisted on corn, wheat, oats, hay and a variety of livestock. Iowa had been converted from a seemingly limitless prairie-forest-wetland mosaic into a domesticated landscape of small farms, grain fields and pastures. There were still undrained sloughs and wet pastures on many farms and tracts of prairie could still be found to remind farmers of vintage Iowa, but these native areas were scattered and becoming ever smaller. In the early 20th century they were still looked on as waste areas needing conversion to a more productive use. Most of Iowa's native wildlife was either gone or reduced to such low numbers that rabbits, squirrels, quail and the occasional prairie chicken were the only game animals available to most hunters.

Change Continues in the 20th Century

The changes in Iowa's landscape in the 20th century were less dramatic but in some ways more devastating. Wildlife and its habitats were impacted by constant improvements in farming technology and the effects of government agricultural policy on farmers' decisions about how their land would be used.

Improved farming technology. Change was slow at first. Much of northern Iowa was too wet to permit iron-wheeled tractors to function so gasoline-powered equipment did not replace horses on a large scale until rubber balloon tires became available in the late 1930's. Hybrid seed corn was introduced in the 1930's to improve yields; for the first time more crop could consistently be raised than was needed for use on the farm. Farming ever so gradually became less a way of life and more of a business.

Industrial technology developed during World War II rapidly accelerated the pace of change. By mid-century mechanical planters, harvesters (hay balers, corn pickers and grain combines) and grain handling equipment were reducing the need for hand labor. Repeated field cultivation for weed control was the norm, but control in cultivated fields was a constant and frequently unsuccessful battle for farmers. Inefficient harvesting equipment often left a substantial part of the crop in the field.

Labor saving devices permitted farmers to handle ever-larger farming operations. In the 1950's the average northern Iowa farm had grown to 250 acres but was still a diverse operation of livestock, small grains, hay and corn. Foxtail-choked cornfields with plenty of waste grain were a pheasant hunter's delight and a source of food and cover for a variety of other game and nongame wildlife.


The last half of the century brought even more change. Modern tiling machines could mechanically dig and insert underground perforated field tiles to drain even the wettest areas. The use of agricultural chemicals – herbicides, pesticides, and fertilizers – became the norm and weeds and insects were, if not conquered, at least minimized as a threat to crop yields. The first pesticides were organochlorines -DDT and its derivatives- that had devastating long-term effects on bird populations that led to the ban on their use in the 1970's. The newer generations of pesticides - organophosphates - have no residual effects that are known at this time, although Roundup, the most widely used herbicide, has recently been implicated in the decline of amphibians.) Soybeans were introduced as a cash crop and genetically modified crops with built-in pesticide resistance were developed. Livestock operations shifted from on-the-farm to confinement operations and the need for extensive livestock forage (hay and small grains) was reduced. Crop rotations eventually were simplified to continuous corn or soybeans or corn-soybean rotations over most of the state. Planting and harvesting equipment and the tractors to pull them became ever larger. Modern grain combines became so efficient that little waste grain or crop residue was left in the fields for wildlife food or cover. By 2000 the average farm had increased to more than 1,000 acres and the number of farms in Iowa had decreased from 150,000 to just 90,000 in a half-century.

USDA farm policies. Government farm policy also played a role in accelerating these changes. Congress passed the first of several programs to retire crop land and spur agricultural income in the depth of the depression in the 1930's. Farm policy shifted to all-out production during World War II. By the mid-1950's farm prices were again depressed and a second, 10-year land retirement program (the Soil Bank) was implemented. Pheasants, bobolinks and other grassland birds responded to the increased habitat until the program ended in 1965.

For the next 20 years USDA required farmers to set aside up to 10 percent of their crop land in order to participate in subsidy programs. These set-aside acres were rotated annually and never developed permanent wildlife cover. Their value to wildlife was limited - some biologists claimed they had a net negative affect on pheasants and other ground-nesting birds because set-aside acres had to be mowed for weed control just at the time birds were nesting.

In the early 1970's grain export quotas were removed to open up international markets. Row crops in Iowa grew by more than 3 million acres at the expense of hay and pasture (Figure 2-1), most in the southern third of Iowa. The distribution of the ringnecked pheasant nearly reversed itself as a result. The new croplands in southern Iowa allowed pheasants to flourish where the bobwhite quail had been the dominant game bird. The added pressure to raise rowcrops eliminated most of the remaining wildlife habitat in northern Iowa, however, and pheasant populations there plummeted.

Figure 2-1. Changes In Cropping Patterns and Representative Grassland Wildlife



The increased row crop acreage also put added pressure on Iowa's remnant forests. Pasture that was converted to row crops had to be replaced, so bulldozing timber to create new pasture became a popular practice. Iowa's forestlands hit their all-time low - 1.5 million acres - during the U.S. Forest Service's 1974 inventory of forestlands.

In the midst of another farm economic crisis in the 1980's a third 10-year land retirement program – the Conservation Reserve Program (CRP) – was introduced to supplement farm income. CRP fields were mostly planted to cool season grasses like smooth brome that provided valuable nesting cover for grassland wildlife. Iowa's pheasant populations and harvest, both in the midst of a 20-year decline, rebounded quickly (Figure 2-2). In northern Iowa pheasant numbers increased wherever CRP fields were planted and increases were also recorded in the southern half of the state. But as the initial 10-year contracts matured the benefits to game birds in southern Iowa declined. Brome developed a thick sod and annual weeds (important foods for birds) were eliminated. Southern Iowa counties that had the maximum of 25 percent of their cropland

Figure 2-2 Statewide Pheasant Trends




Figure 2. Mean number of pheasants counted on 30-mile August roadside survey routes, statewide, 1962-present compared to total statewide pheasant harvest.

enrolled in CRP saw declines in pheasants and quail. IDNR-sponsored research would eventually find that nongame birds like Henslow's sparrow that nested in mature grasslands would respond to the habitat provided by older CRP fields.


Small mammals and the avian and mammalian predators that fed on them would increase also. The return of the bobcat to Iowa is at least partly explained by the prey provided in CRP fields.

CRP acreages in whole fields peaked at 2.2 million acres, but modifications in the late 1990's and early 21st century reduced whole-field enrollments to 1.5 million acres today. Most contracts will expire in 2007 and 2008. Recent farm bills have included a number of permanent and short-term programs designed to provide soil and wildlife conservation benefits as well as subsidize the production of commodity crops. The Continuous CRP (buffer strips), Wetland Reserve Program (WRP), Wildlife Habitat Incentive Program (WHIP), Farmed Wetland Program (FWP) and others have been beneficial, but most have been implemented on smaller parcels than the original CRP fields. Potential problems with habitat fragmentation, connectivity between habitat blocks and their value to area-sensitive species is not well understood (see Chapter 5).

Summary. The result of this improved technology and the flurry of often-conflicting farm legislation has been a gradual and long-term decline in wildlife habitat on private agricultural lands. Farm operations have shifted from diversified agriculture to corn and soybean monocultures. Between 1900 and 2004 row crop acreages increased from 9.1 million acres to 21 million acres (Figure 2-1). Hay and small grain acreage decreased from 6.8 million acres to a current 1.1 million acres. Larger farms and field sizes have eliminated fencerows, windbreaks, waterways and other on-farm habitat. The nearly exclusive use of farm chemicals for weed and insect control has eliminated food and cover for songbirds and other wildlife. Conservation practices subsidized by various titles of recent farm legislation have helped slow this trend, but the funding available to implement them has never equaled the amount USDA has spent subsidizing commodity crops that encourages maximum production.

The impact on of these trends on wildlife that utilizes agricultural lands has been slowly devastating and is the subject of much of the remainder of this Plan. The loss of grasslands as row crop agriculture has become dominant has resulted in substantial declines in bobolinks and white-tailed jackrabbits (Figure 2-1), two once-common native grassland species. Even the popular ringnecked pheasant, until recently the state's most well known game animal (Figure 2-2) is in the midst of a 50-year decline in numbers. A similar trend can be seen for bobwhite quail (Figure 2-3). Other examples can be found in *Trends in Iowa Wildlife Populations and Harvest* (2004 and earlier years) published by IDNR.

Figure 2-3. Trends In Bobwhite Quail Populations and Harvest.



These landscape changes have impacted aquatic wildlife as well, although they are not as well documented. Advertisements to attract settlers to Iowa in the 1850's stressed the vast acreages of fertile soils, abundant wildlife and sparkling clean waters teeming with game fish. A muskellunge museum specimen at Iowa State University was caught on the Skunk River near Ames in the 1880's.

By the early 20th century, however, conservationists Aldo Leopold and Jay N, "Ding" Darling were decrying the excessive erosion of soils that had been denuded of their vegetative cover and the excessive siltation of Iowa's waters that resulted. Loss of vegetative cover, excessive grazing, channelization of streams, and shoreline alterations led to accelerated siltation and the transport of pesticides and fertilizers into aquatic systems from agricultural fields. Heavy silt loads altered water turbidity and temperature regimes. Streambed degradation and the loss of submergent and emergent plants frequently followed. As the silt settles it can cover existing bottom substrates and alter the entire natural community.

All of these alterations to native habitats, aquatic plant communities and wildlife increase the opportunities for invasive exotic species to supplant native wildlife. Alien species like carp further increased water turbidity and in many cases made smaller water bodies unsuitable for native fish.

Wildlife Conservation

Wildlife Restoration. Not all wildlife trends of the past half-century have been negative. The creation of the Iowa State Conservation Commission (now IDNR) in 1935, the gradual development of wildlife science and management as professions after World War II, and the formation of IDNR's Wildlife Diversity Program in 1981 have returned a portion of Iowa's native wildlife to the state. White-tailed deer, wild turkeys and giant Canada geese are now more abundant than at any time since the late 1800's. Other restoration programs have returned ruffed grouse to southern Iowa, river otters to the state's streams, peregrine falcons, ospreys and trumpeter swans nest again in Iowa. Bald eagles, bobcats and Sandhill cranes have reappeared as a result of successful conservation programs elsewhere. Details of these and other wildlife restoration programs are explained in *Trends in Iowa Wildlife Populations and Harvest - 2004*.

Land acquisition. IDNR has also pursued land acquisition programs to permanently protect and enhance wildlife habitat. Since 1972 Iowa waterfowlers have been required to purchase an Iowa Migratory Game Bird Stamp. Since 1979 all hunters have been required to purchase a Habitat Stamp along with their hunting license. Proceeds from these stamps are dedicated to habitat protection and management. Funds from the Habitat Stamp are shared equally with Iowa's 99 County Conservation Boards.


IDNR has aggressively sought funds for habitat protection through the North American Waterfowl Management Plan, the North American Wetlands Conservation Act, State Wildlife Grants, the Environmental Protection Agency, Iowa County Conservation Boards and others. IDNR also partners with a number of NGOs to extend the reach of state and Federal funds. Ducks Unlimited, Pheasants Forever, National Wild Turkey Federation, Iowa Natural Heritage Foundation and The Nature Conservancy have been major cooperators with IDNR's habitat protection programs. Numerous other NGO's and individual private contributor have helped as well.

Since 1980 IDNR has acquired 100,000 acres of land to enhance or restore habitat for wildlife. Research sponsored by IDNR has found that these restored lands are colonized quickly by birds that utilize the appropriate habitats as they develop. Little is known about responses from less-mobile wildlife like amphibians and reptiles.

In spite of the aggressive efforts to protect wildlife habitat, Iowa remains one of the states with the highest proportion of privately held land (Map 2-2). Public conservation lands accounted for just over 600,000 acres in 2004, or just 1.7% of the land area of the state (Iowa GAP).

The IDNR owns nearly half of the public conservation lands (322,00 acres), including state parks, state forests and wildlife management areas. Federal ownership accounts for 138,000 acres in four flood control reservoirs, 5 national wildlife refuges and 57,000 acres of WRP easements. IDNR has land management agreements on portions of the reservoirs but little control over water levels. County Conservation Boards own 134,000 acres.

Map 2-2. Publicly-owned Conservation Lands in Iowa



Habitat on private lands. Wildlife habitats on private lands have also received attention from IDNR programs. Farm Game Habitat crews roamed the state in the 1950's and 1960's helping landowners establish habitat on their property. In 1971 the number of IDNR wildlife management biologists was doubled and they were housed in USDA farm service center offices to promote contacts with private landowners. In the 1980's farmstead shelterbelts and switchgrass cost-sharing programs were introduced to promote these practices on private land. For the past 20 years IDNR biologists have actively promoted USDA farm bill practices (e.g. CRP, WRP) that provide landowners funds to assist with developing wildlife t. The Wildlife Bureau's Private lands Program was formed in 2002 to take better advantage of wildlife-friendly USDA farm programs and other Federal grants like the LIP.

The success of private lands assistance programs has been mixed. Wildlife habitat has been affected on 65,000 acres. In the past, however, private land habitats have disappeared when land ownership or landowner objectives change or as government subsidies for habitat protection have ended. Some of the new programs (WRP, LIP) require permanent easements and habitat improvements will remain. Others are short term and will likely revert to more financially rewarding uses when government programs end.


Iowa's Natural Communities Today

The result of a century and a half of change as a result of human intervention on Iowa's landscape has been a shift in the composition of Iowa's plant communities and the wildlife that inhabits them. Few undisturbed natural plant or wildlife communities exist today. Less than 0.1% of Iowa's native prairies (30,000 acres), 5% of its wetlands (422,000 acres), and 43% of its forests (2,800,000 acres) remain.

Of these, forestlands are the only habitats that are increasing on private lands without state or Federal intervention. The shift from raising livestock on the farm to confinement operations has reduced the need for pasture in the past decade and a half. Today Iowa has nearly 1 million acres of pastureland reverting to early successional forest, most in the Southern Iowa Drift Plain, the Paleozoic Plateau and the Loess Hills.

Map 2-3 shows the land cover in Iowa in the year 2002. The majority of the state is covered with row crop, primarily corn and soybeans. Most of the remainder of the state is in grassland, often conservation reserve, road ditches or pasture, with lesser acreages of timber and other habitat types. More details on the current status of Iowa's wildlife are provided in Chapter 3, and the status of wildlife habitats in Chapter 4.

Map 2-3. Landcover of Iowa in 2002 (IDNR Landsat 5 and 7 Enhanced Thematic Mapper Images)



CHAPTER THREE

SPECIES OF GREATEST CONSERVATION NEED

Wildlife Species Considered by the IWAP

Federal guidelines require the IWAP to include information on the distribution and abundance of wildlife, including low and declining populations *as the IDNR deems appropriate*. The IDNR is the sole agency given the responsibility to manage Iowa's fish and wildlife resources and to protect their habitats (Code of Iowa, Chapter 455A). Iowa law defines *wildlife* as any species of wild mammal, fish, bird, reptile and amphibian (Code of Iowa sections 456.24, 481A.1, 481A.38, 481A.39, 481A.48). Authority to establish and protect state-listed endangered or threatened species is vested in Chapter 481B.4 and Iowa Administrative Code Chapter 571-77(481B). Butterflies, land snails and fresh water mussels were included in the Plan because these invertebrates are included on the state's endangered and threatened species list. Dragonflies and damselflies were later added when significant data were found that listed the distribution and status of species in these groups. A total of 999 species were considered by working groups (Table 3-1).

Table 3-1. Number of species considered by the IWAP.

Taxonomic Class	Species	List location
Breeding birds	206	Appendix 3
Migrant birds ¹	199	Appendix 4
All birds	405	
Mammals	88	Appendix 5
Fish	153	Appendix 6
Reptiles and Amphibians	71	Appendix 7
Freshwater mussels	55	Appendix 8
Land snails	8	Table 3-8
Butterflies	113	Appendix 9
Dragonflies and Damselflies	106	Appendix 10
Total species considered	999	

¹ Migrant species that do not nest in Iowa

Determining the Species of Greatest Conservation Need

A variety of data resources were utilized by working groups as they selected the SGCN:

- Iowa GAP -completed in 2003 with ongoing updates provided by Iowa Nature Mapping;

- Published historic and scientific literature;
- Unpublished reports, scientific surveys and databases maintained by the IDNR fisheries, wildlife and water quality bureaus;
- Personal research and survey data supplied by wildlife ecologists at Iowa educational institutions;
- Museum and personal specimen collections;
- State and regional databases maintained by other conservation organizations (e.g. NatureServe, PIF, PARC, TNC, USFWS, IOU, Audubon IBA, etc.);
- Personal expertise of working group members and consultants.

The procedures used to identify SGCN varied somewhat between Working Groups, but all followed a similar process. Working Group members reviewed the available information on species abundance and distribution in their assigned taxonomic class. The potential list of SGCN in each taxonomic class was then circulated to other experts for review and comment. The Working Group reviewed comments and prepared a recommended list that was presented to the Steering Committee for approval. In some cases additional species were added to the lists based on comments received later in the planning process.

Game species. Information on the current abundance and distribution of wildlife is most complete for major game species. IDNR has collected harvest statistics and conducted population surveys on some of these species for over half a century. This information is documented for terrestrial game and selected nongame species in *Trends in Iowa Wildlife Populations and Harvest 2004* (Iowa DNR 2004). *Fishing in Iowa: A Survey of 1994 Iowa Anglers* (Iowa DNR 1995) is the most recent survey of statewide fish harvest statistics. A compilation of population surveys is provided in Appendix 21 and summarized in Table 7-1.

Birds. The distribution and abundance of birds in Iowa is better understood than any other nongame taxa considered in the IWAP. As a result the Bird Working Group had many sources of information to consult. The working group used the bird list of the IOU to create separate state and national status lists for all nesting or migrant species that are found in Iowa. Other Iowa and national status lists were created using the NatureServe website, the NABCI 2002 Bird Conservation Region and the USFWS Region 3 Birds of Conservation Concern lists. Once completed, these lists were compared, reviewed by the working group and collated into final lists of Iowa breeding and migrant birds that were accepted by the Steering Committee.

To determine the bird SGCN, a subcommittee of the Bird Working Group classified breeding and migrant birds into one of 3 categories:

- 1 - SGCN (State Endangered, Threatened, or Special Concern);
- 2 - Strong candidate for SGCN;
- 3- Questionable candidate for SGCN.

This list was provided to the entire Bird Working Group for review. Then a second draft of was created in which birds were listed in one of two categories:

- 1= SGCN (State Endangered, Threatened, and Special Concern);
- 2 = Proposed for SGCN by the Bird Working Group.

This second draft was reviewed by the entire working group and USFWS Region 3 biologists, resulting in a final list of 67 Iowa breeding birds (Table 3-2) and 18 migratory birds (Table 3-3) of greatest conservation need. Iowa has a responsibility to conserve these migrant species due to the state's geographical location on their migratory pathway, (e.g. rusty blackbird, greater and lesser yellowlegs), or because Iowa acts as important migration rest habitat (e.g. yellow rail).

There are several Iowa extirpated nesting bird species not included as birds of greatest conservation need e.g. swallow-tailed kite (*Elanoides forficatus*), merlin (*Falco columbarius*), and common loon (*Gavia immer*). Restoration programs for these species may be justified in the future, at which time their status will be reconsidered.

The Partners in Flight (PIF) Species Prioritization Scheme (1991) played a major role in determining which species ultimately made the bird list of SGCN. A number of the priority bird species lists used as references in this process were derived directly or indirectly from PIF. The PIF prioritizing system ranks each species of North American breeding bird based upon seven measures of conservation vulnerability:

- 1) Relative abundance (inter-specific);
- 2) Size of breeding range;
- 3) Size of non-breeding range;
- 4) Threats to the species in breeding areas;
- 5) Threats to the species in non-breeding areas;
- 6) Population trend;
- 7) Relative density (intra-specific) in a given planning unit compared to the maximum reached within its range.

Additional references utilized by the Bird Working Group include:

- NatureServe National and Sub-national Heritage Status Rankings;
- *Iowa Birds* (Dinsmore et. al.1984);
- *The Iowa Breeding Bird Atlas* (Jackson et. al. 1996);
- *Birds of Iowa* (Kent and Dinsmore 1996);
- USGS Breeding Bird Survey;
- Partners in Flight Bird Conservation Plans for Iowa Physiographic Areas;
- NABCI 2002 Bird Conservation Region
- USFWS Region 3 Birds of Conservation Concern;
- United States Shorebird Conservation Plan;
- Upper Mississippi-Great Lakes Shorebird Plan;
- Northern Plains/Prairie Potholes Regional Shorebird Plan;

- North American Waterbird Conservation Plan;
- The Prairie-Forest Border Ecoregion: A Conservation Plan (The Nature Conservancy);
- American Bird Conservancy Green List;
- Iowa Important Bird Area Priority Birds List (Audubon).

Mammals. The Mammal Working Group developed a complete list of mammals found in Iowa primarily based on Bowles *et al.* 1998. The SGCN list for mammals was developed by eliminating vagrant and extirpated species, which had little possibility for management. The working group then eliminated all species that had a NatureServe S + N status score of 9 or above (*secure* or *apparently secure*) at the state and national levels (Appendix 11). The proposed SGCN list was sent to individuals outside of the working group for comment and was reviewed by the entire Steering Committee. The final list of SGCN is comprised of 18 species (Table 3-4).

Amphibians and reptiles. This working group developed a complete list of Iowa amphibians and reptiles based primarily the work of Christiansen and Bailey (1986, 1988, and 1991). Those species with a NatureServe S plus N status score of 9 or above (*secure* or *apparently secure*) were eliminated if they were not on Iowa's T & E list or did not have a NatureServe trend listing of declining or unknown. The final list of amphibians and reptiles of greatest conservation need contains 31 species (Table 3-5).

Land snails. Comparatively little is known about the distribution and status of this group in Iowa and there is no comprehensive list of land snails for the state. For this reason, the Land Snails Working Group decided to list only those species on the state's T & E list (Appendix 12). This list was based largely on comprehensive surveys of algific slopes in NE Iowa conducted by Dr. Terrence Frest in the 1980's. Eight species were listed as Iowa land snails of greatest conservation need (Table 3-6).

Butterflies. The Butterfly Working Group first developed a complete list of Iowa butterflies (Appendix 9). Knowledge of butterfly abundance and distribution is not as complete as for other taxonomic groups considered in the Plan, so only Iowa listed T & E species were included on the SGCN list for butterflies. Both lists were reviewed by the working group and by the full Steering Committee. The final list of Iowa butterflies of greatest conservation need contains 30 species (Table 3-7).

Table 3-2. Breeding birds of greatest conservation need.

Iowa Abundance: A = abundant, C = common, CL = common locally, U=uncommon, UL = uncommon locally, R = rare, SC = special concern, Th = threatened, En = endangered, X = extirpated. E = extinct.

Iowa Trend: D = decreasing, I = increasing, S = stable, K = unknown.

Iowa Status: B =breeding, N = non-breeding.

See Appendix 11for NatureServe codes used in Iowa Status and National Status columns.

Common Name	Scientific Name	Iowa Abundance	Iowa Trend	Iowa Status	National Status
American bittern	<i>Botaurus lentiginosus</i>	R	S	S2B	N4B, N4N
Least bittern	<i>Ixobrychus exilis</i>	U	S	S3B, S2N	N5B, N5N
Black-crowned night-heron	<i>Nycticorax nycticorax</i>	R	D	S3B, S3N	N5B, N5N
Yellow-crowned night-heron	<i>Nyctanassa violacea</i>	R	K	S3B, S3N	N5B, N5N
Trumpeter swan	<i>Cygnus buccinator</i>	R	I	S2B	N4B, N4N
Northern pintail	<i>Anas acuta</i>	R	S	S2B, S5N	N5B, N5N
Canvasback	<i>Aythya valisineria</i>	R	S	S2B, S4N	N5B, N5N
Redhead	<i>Aythya americana</i>	U	S	S2B, S4N	N5B, N5N
Osprey	<i>Pandion haliaetus</i>	R	I	SXC, S3N	N5B, N4N
Bald eagle	<i>Haliaeetus leucocephalus</i>	En	I	S3B, S3N	N4B, N4N
Northern harrier	<i>Circus cyaneus</i>	En	I	S2B, S4N	N5B, N5N
Red-shouldered hawk	<i>Buteo lineatus</i>	En	I	S2B	N5B, N5N
Broad-winged hawk	<i>Buteo platypterus</i>	R	S	S3B	N5B
Swainson's hawk	<i>Buteo swainsoni</i>	R	D	S3B, S3N	N5B
Peregrine falcon	<i>Falco peregrinus</i>	En	I	S1B	N4B, N4N
Ruffed grouse	<i>Bonasa umbellus</i>	U	D	S4B	N5
Greater prairie-chicken	<i>Tympanuchus cupido</i>	R	S	S1B	N4
Sharp-tailed Grouse	<i>Tympanuchus phasianellus</i>	R	K	S1B	N4
Northern bobwhite	<i>Colinus virginianus</i>	CL	D	S5B	N5
King rail	<i>Rallus elegans</i>	En	K	SAB, S1N	N4B, N4N
Common moorhen	<i>Gallinula chloropus</i>	R	K	S2B, S2N	N5B, N5N
Sandhill crane	<i>Grus canadensis</i>	R	I	S1B, S1N	N5B, N5N
Piping plover	<i>Charadrius melodus</i>	En	S	S1B	N3B, N3N
Upland sandpiper	<i>Bartramia longicauda</i>	U	S	S3B	N5B
American woodcock	<i>Scolopax minor</i>	C	K	S4B, S5N	N5B, N5N
Wilson's phalarope	<i>Phalaropus tricolor</i>	R	K	S3N	N5B
Forster's tern	<i>Sterna forsteri</i>	SC	K	S2B, S3N	N5B, N5N
Least tern	<i>Sterna antillarum</i>	En	S	S1B, S1N	N?
Black tern	<i>Chlidonias niger</i>	SC	D	S1B, S4N	N4B
Black-billed cuckoo	<i>Coccyzus erythrophthalmus</i>	C	D	S3B	N5B

Common Name	Scientific Name	Iowa Abundance	Iowa Trend	Iowa Status	National Status
Yellow-billed cuckoo	<i>Coccyzus americanus</i>	C	D	S3B	N5B
Barn owl	<i>Tyto alba</i>	En	S	S1B	N5
Burrowing owl	<i>Speotyto cunicularia</i>	R	K	S1B	N4B, N4N
Long-eared owl	<i>Asio otus</i>	Th	K	S2B, S3N	N5B, N5N
Short-eared owl	<i>Asio flammeus</i>	En	S	S1B, S2N	N5B, N5N
Common nighthawk	<i>Chordeiles minor</i>	C	D	S5B	N5B
Whip-poor-will	<i>Caprimulgus vociferus</i>	CL	D	S5B	N5B
Red-headed woodpecker	<i>Melanerpes erythrocephalus</i>	C	D	S5B	N5B, N5N
Acadian flycatcher	<i>Empidonax virescens</i>	R	S	S3B, S3N	N5B
Willow flycatcher	<i>Empidonax traillii</i>	C	S	S4B, S4N	N5B
Least flycatcher	<i>Empidonax minimus</i>	R	K	S1B, S4N	N5B
Brown creeper	<i>Certhia americana</i>	R	S	S3B	N5
Bewick's wren	<i>Thryomanes bewickii</i>	R	K	S2B, S2N	N5B
Sedge wren	<i>Cistothorus platensis</i>	U	I	S4B, S4N	N4B, N5N
Veery	<i>Catharus fuscescens</i>	R	D	S2B, S3N	N5B
Wood thrush	<i>Hylocichla mustelina</i>	U	D	S4B, S4N	N5B
Northern mockingbird	<i>Mimus polyglottos</i>	R	K	S3B	N5
Loggerhead shrike	<i>Lanius ludovicianus</i>	U	D	S3B, S3N	N4
White-eyed vireo	<i>Vireo griseus</i>	R	K	S2B, S3N	N5B, N5N
Bell's vireo	<i>Vireo bellii</i>	U	S	S3B, S4N	N4B
Blue-winged warbler	<i>Vermivora pinus</i>	R-W/U-E	S	S3B, S4N	N5B
Cerulean warbler	<i>Dendroica cerulea</i>	R	D	S2B, S3N	N4B
Black-and-white warbler	<i>Mniotilta varia</i>	R	K	S5N	N5B, N4N
Prothonotary warbler	<i>Protonotaria citrea</i>	R	S	S3B, S3N	N5B
Worm-eating warbler	<i>Helmitheros vermivorus</i>	R	K	S2B, S2N	N5B
Louisiana waterthrush	<i>Seiurus motacilla</i>	R	K	S3B, S4N	N5B
Kentucky warbler	<i>Oporornis formosus</i>	R	K	S1B, S3N	N5B
Hooded warbler	<i>Wilsonia citrina</i>	R	K	S1B, S2N	N5B
Yellow-breasted chat	<i>Icteria virens</i>	R	S	S3B, S3N	N5B
Dickcissel	<i>Spiza americana</i>	A	D	S4B, S4N	N5B
Eastern towhee	<i>Pipilo erythrophthalmus</i>	CL	D	S4B, S4N	N5
Field sparrow	<i>Spizella pusilla</i>	C	D	S5B, S5N	N5
Lark sparrow	<i>Chondestes grammacus</i>	CL	K	S4B	N5B
Grasshopper sparrow	<i>Ammodramus savannarum</i>	C	D	S4B, S4N	N5B, N5N
Henslow's sparrow	<i>Ammodramus henslowii</i>	Th	I	S3B, S2N	N3B, N4N

Common Name	Scientific Name	Iowa Abundance	Iowa Trend	Iowa Status	National Status
Bobolink	<i>Dolichonyx oryzivorus</i>	C	D	S4B	N5B
Eastern meadowlark	<i>Sturnella magna</i>	C	D	S4B, S4N	N5

Table 3-3. Migratory birds of greatest conservation need.

Iowa Abundance: R = rare, U = uncommon, C = common

See Appendix 11 for NatureServe codes used in Iowa Status and National Status columns.

Common Name	Scientific Name	Iowa Abundance	Iowa Status	National Status
American white pelican	<i>Pelecanus erythrorhynchos</i>	C	S4N	N3B, N3N
Yellow rail	<i>Coturnicops noveboracensis</i>	R	SNA	N3B, N4N
Whooping crane	<i>Grus americana</i>	R	SXB	N1N
American golden-plover	<i>Pluvialis dominica</i>	U	S5N	N5B, N5N
American avocet	<i>Recurvirostra americana</i>	R	S3N	N5B, N5N
Greater yellowlegs	<i>Tringa melanoleuca</i>	C	S5N	N5B, N5N
Lesser yellowlegs	<i>Tringa flavipes</i>	C	S5N	N5B, N5N
Solitary sandpiper	<i>Tringa solitaria</i>	C	S5N	N4B, N5N
Hudsonian godwit	<i>Limosa haemastica</i>	U	S3N	N3?B
Marbled godwit	<i>Limosa fedoa</i>	R	SXB	N5B, N5N
Stilt sandpiper	<i>Micropalama himantopus</i>	U	S4N	N3B, N4N
Buff-breasted sandpiper	<i>Tryngites subruficollis</i>	R	S3N	N4B
Short-billed dowitcher	<i>Limnodromus griseus</i>	U	S4N	N5B, N5N
Golden-winged warbler	<i>Vermivora chrysoptera</i>	U	S1N	N4B
Canada warbler	<i>Wilsonia canadensis</i>	U	S3N	N5B
Le Conte's sparrow	<i>Ammodramus leconteii</i>	U	SNA	N3B, N4N
Nelson's sharp-tailed sparrow	<i>Ammodramus nelsoni</i>	R	SNA	N3B, N5N
Rusty blackbird	<i>Euphagus carolinus</i>	U	S3N	N5B, N5N

Table 3-4. Mammals of greatest conservation need.

Iowa Abundance: CL = common locally, U = uncommon, R = rare,
I = re-introduced.

Iowa Trend: K = unknown, D = decreasing, S = stable, I = increasing.

See Appendix 11 for NatureServe codes used in Iowa Status and National Status columns

Common Name	Scientific Name	Iowa Abundance	Iowa Trend	Iowa Status	National Status
Hayden's shrew	<i>Sorex haydeni</i>	CL	K	S4-Apparently secure	N4
Short-tailed shrew	<i>Blarina hylophaga</i>	CL	K	S4	?
Least shrew	<i>Cryptotis parva</i>	R	K	S3 Threatened	N5
Evening bat	<i>Nycticeius humeralis</i>	CL	K	S3	N5
Indiana bat	<i>Myotis sodalis</i>	R	K	S1 Endangered	N2
Northern myotis	<i>Myotis septentrionalis</i>	CL	K	S4-Apparently secure	N4
White-tailed jackrabbit	<i>Lepus townsendii</i>	R	D	S3	N5
Franklin's ground squirrel	<i>Spermophilus franklinii</i>	R	D	S3	N5
Red squirrel	<i>Tamiasciurus hudsonicus</i>	CL	S	S3	N5
Southern Flying Squirrel	<i>Glaucomys volans</i>	U	K	S4 Special Concern	N5
Plains pocket mouse	<i>Perognathus flavescens</i>	R	K	S2	N5
Prairie vole	<i>Microtus ochrogaster</i>	U	S	S3	N5
Red-backed vole	<i>Clethrionomys gapperi</i>	R	D	S2 Endangered	N5
Southern bog lemming	<i>Synaptomys cooperi</i>	R	D	S3 Threatened	N5
Woodland vole	<i>Microtus pinetorum</i>	R	K	S3	N5
River otter	<i>Lutra canadensis</i>	I	I	S3	N5
Spotted skunk	<i>Spilogale putorius</i>	R	D	S1 Endangered	N5
Bobcat	<i>Lynx rufus</i>	U	I	S3	N5

Table 3-5. Reptiles and amphibians of greatest conservation need.

Iowa Listing: T = Threatened, E = Endangered

Iowa Trend: K = unknown, D = decreasing, S = stable.

See Appendix 11. for NatureServe codes used in Iowa Status and National Status columns

Common Name	Scientific Name	Iowa Listing	Iowa Trend	Iowa Status	National Status
Mudpuppy	<i>Necturus maculosus</i>	T	D	S2	N5
Central Newt	<i>Notophthalmus viridescens</i>	T	S	S2	N5
Smallmouth Salamander	<i>Ambystoma texanum</i>		S	S3	N5
Blue-spotted Salamander	<i>Ambystoma laterale</i>	E	S	S1	N5
Crawfish Frog	<i>Rana areolata</i>	E	D	S1	N4
Cricket Frog	<i>Acris crepitans</i>		D	S3	N5
Great Plains Toad	<i>Bufo cognatus</i>		D	S3	N5
Ornate Box Turtle	<i>Terrapene ornata</i>	T	S	S2	N5
Blanding' Turtle	<i>Emydoidea blandingii</i>	T	D	S2	N4
Wood Turtle	<i>Clemmys insculpta</i>	E	K	S1	N4
Alligator Snapping Turtle	<i>Macrochelys temmincki</i>		K	SU	N3, N4
Yellow Mud Turtle	<i>Kinosternon flavescens</i>	E	D	S1	N5
Common Musk Turtle	<i>Sternotherus odoratus</i>	T	D	S2	N5
Slender Glass Lizard	<i>Ophisaurus attenuatus</i>	T	D	S2	N5
Six-Lined Racerunner	<i>Cnemidophorus sexlineatus</i>		D	S3	N5
Northern Prairie Skink	<i>Eumeces septentrionalis</i>		D	S3	N5
Great Plains Skink	<i>Eumeces obsoletus</i>	E	D	S1	N5
Diamondback Water Snake	<i>Nerodia rhombifera</i>	T	D	S2	N5
Yellowbelly Water Snake	<i>Nerodia erythrogaster flavigaster</i>		D	S1	N5
Copperbelly Water Snake	<i>Nerodia erythrogaster neglecta</i>	E	D	S1	N5
Smooth Earth Snake	<i>Virginia valeriae</i>		S	S3	N5
Western Worm Snake	<i>Carphophis amoenus</i>	T	D	S2	N5
Smooth Green Snake	<i>Opheodrys vernalis</i>	Special Concern	S	S3	N5
Prairie Kingsnake	<i>Lampropeltis calligaster</i>		D	S3	N5
Speckled Kingsnake	<i>Lampropeltis getulus</i>	T	D	S1	N5
Bullsnake	<i>Pituophis catenifer sayi</i>	Special Concern	D	S3	N5
Western Hognose Snake	<i>Heterodon nasicus</i>	E	K	S1	N5
Eastern Massasauga Rattlesnake	<i>Sistrurus catenatus catenatus</i>	E (Fed. Candidate Sp.)	D	S1	N3, N4
Timber Rattlesnake	<i>Crotalus horridus</i>		D	S3	N5
Prairie Rattlesnake	<i>Crotalus viridis</i>	E	K	S1	N5
Copperhead	<i>Agkistrodon contortix</i>	E	D	S1	N5

Table 3-6. Land snails of greatest conservation need.

Iowa and Federal Listing: T = T, E = E.

Iowa Abundance: R = Rare

Iowa Trend: K = unknown, S = stable, D = decreasing.

See Appendix 11 for NatureServe codes used in Iowa Status and National Status columns

Common Name	Scientific Name	Iowa Listing	Federal Listing	Iowa Abundance	Iowa Trend	National Status
Iowa Pleistocene Snail	<i>Discus macclintocki</i>	E	E	R	S	N1
Frigid Ambersnail	<i>Catinella gelida</i>	E		R	D	N1
Minnesota Pleistocene Succinea	<i>Novasuccinea n. Sp. Minnesota a</i>	E		R	D	NNR
Iowa Pleistocene Succinea	<i>Novasuccinea n. Sp. Minnesota b</i>	E		R	D	NNR
Briarton Pleistocene Snail	<i>Vertigo brierensis</i>	E		R	K	N1
Hubricht's Vertigo	<i>Vertigo hubrichti</i>	T		R	K	N3
Iowa Pleistocene Vertigo	<i>Vertigo iowaensis</i>	E		R	K	N3
Bluff Vertigo	<i>Vertigo occulta</i>	T		R	K	N2

Table 3-7. Butterflies of greatest conservation need.

Iowa Listing and Abundance: K = unknown, R = rare, C = common, LC = locally common.

Iowa Abundance: K = unknown, S = stable, D = decreasing.

See Appendix 11 for NatureServe codes used in Iowa Status and National Status columns

Common Name	Scientific Name	Iowa Listing	Iowa Abundance	Iowa Trend	Iowa Status	National Status
Pepper and Salt Skipper	<i>Amblyscirtes hegon</i>	SC	K	K	S?	N5
Arogos Skipper	<i>Atrytone arogos</i>	SC	R	D	S2	N3
Dusted Skipper	<i>Atrytonopsis hianna</i>	SC	R	D	S3	N4
Pipevine Swallowtail	<i>Battus philenor</i>	SC	K	K	S?	N5
Swamp Metalmark	<i>Calephelis muticum</i>	SC	R	K	S?	N3
Common Ringlet	<i>Coenonympha tullia</i>	E	R	D	S1	N5
Wild Indigo duskywing	<i>Erynnis baptisiae</i>	SC	K	K	S3	N5
Sleepy Duskywing	<i>Erynnis brizo</i>	SC	R	K	S3	N5
Dreamy Duskywing	<i>Erynnis icelus</i>	SC	R	K	S3	N5
Columbine Duskywing	<i>Erynnis lucilius</i>	SC	R	K	S3	N4
Olympia White	<i>Euchloe olympia</i>	SC	K	K	S3	N4
Baltimore Checkerspot	<i>Euphydryas phaeton</i>	T	R	K	S2	N4
Two-spotted Skipper	<i>Euphyes bimacula</i>	SC	R	K	S2	N4
Sedge Skipper	<i>Euphyes dion</i>	SC	K	K	S3	N4
Zebra Swallowtail	<i>Eurytides marcellus</i>	SC	R	K	S?	N5
Silvery Blue	<i>Glaucopsyche lygdamus</i>	T	R	K	S2	N5
Dakota Skipper	<i>Hesperia dacotae</i>	E	R	D	S1	N2
Leonardus Skipper	<i>Hesperia leonardus</i>	SC	R	D	S2	N4
Ottoe Skipper	<i>Hesperia ottoe</i>	SC	R	D	S2	N3
Purplish Copper	<i>Lycaena helloides</i>	SC	R	K	S?	N5
Powesheik Skipperling	<i>Oarisma powesheik</i>	T	R	D	S2	N2
Mulberry Wing	<i>Poanes massasoit</i>	T	R	D	S2	N4

Common Name	Scientific Name	Iowa Listing	Iowa Abundance	Iowa Trend	Iowa Status	National Status
Broad-winged Skipper	<i>Poanes viator</i>	SC	LC	K	S3	N5
Zabulon Skipper	<i>Poanes zabulon</i>	SC	LC	K	S3	N5
Byssus Skipper	<i>Problema byssus</i>	T	R	D	S2	N3
Acadian Hairstreak	<i>Satyrium acadica</i>	SC	LC	S	S3	N5
Hickory Hairstreak	<i>Satyrium caryaevorum</i>	SC	R	K	S3	N4
Edward's Hairstreak	<i>Satyrium edwardsii</i>	SC	K	K	S3	N5
Striped Hairstreak	<i>Satyrium liparops</i>	SC	K	K	S?	N5
Regal Fritillary	<i>Speyeria idalia</i>	SC	K	K	S2	N3

Fish. The Fish Working Group first developed a complete list of 153 species of fish found in Iowa and the border rivers based primarily on Harlan and Speaker (1987). Species were placed on a list of greatest conservation need if they were Federal or state listed T and E species; species with recent, rapid and or widespread decrease in abundance or distribution; or species that are targets of regional or national conservation projects. Species with an S plus N NatureServe status score of 8 or greater (*secure* or *apparently secure*) were also eliminated. This list was further refined through comments from IDNR fisheries supervisors and biologists. The complete list was ultimately reduced to 68 fish species of greatest conservation need (Table 3-8).

Freshwater mussels. The complete list of Iowa freshwater mussels (Appendix 6) was based primarily on Arbuckle and Downing (2000) and Heidebrink (2002). Since most Iowa mussel species are decreasing in abundance, additional information on regional abundance, trend, and listing was used to determine species of greatest conservation need. A list of SGCN was developed by eliminating species with an S plus N NatureServe status score of 9 or greater and species that are considered extirpated from this state (unless the species was on the T & E list). This list was reviewed by the working group and the Steering Committee. The final list of mussels of greatest conservation need contains 29 species (Table 3-9).

Table 3-8. Fish of greatest conservation need.

Iowa Abundance: A = abundant, C = common, U = uncommon, R = rare, K = unknown, X = possibly extirpated.

Iowa Trend: K = unknown, I = increasing, D = decreasing, S = stable.

See Appendix 11 for NatureServe codes used in Iowa Status and National Status columns

Common Name	Scientific Name	Iowa Abundance	Iowa Trend	Iowa Status	National Status
Chestnut lamprey	<i>Ichthyomyzon castaneus</i>	R	K	S2	N4
Northern brook lamprey	<i>Ichthyomyzon fossor</i>	R	K	S3	N4
Silver lamprey	<i>Ichthyomyzon unicuspis</i>	U	K	S3	N5
American brook lamprey	<i>Lampetra appendix</i>	C	K	S3	N4
Lake sturgeon	<i>Acipenser fulvescens</i>	R	K	S1	N3N4
Pallid sturgeon	<i>Scaphirhynchus albus</i>	R	D	S1	N1
Shovelnose sturgeon	<i>Scaphirhynchus platyrhynchus</i>	C	S	S4	N4
Paddlefish	<i>Polyodon spathula</i>	C	S	S3	N4
Bowfin	<i>Amia calva</i>	C	S	S3	N5
Longnose gar	<i>Lepisosteus osseus</i>	C	S	S3	N5
Spotted gar	<i>Lepisosteus oculatus</i>	K	K	S3	N5
American eel	<i>Anguilla rostrata</i>	R	D	S2	N5
Skipjack herring	<i>Alosa chrysochloris</i>	U	K	S3	N5
Goldeye	<i>Hiodon alosoides</i>	U	K	S3	N5
Brook Trout	<i>Salvelinus fontinalis</i>	U	S	S3	N5
Grass pickerel	<i>Esox americanus</i>	U	S	S3	N5
Central mudminnow	<i>Umbra limi</i>	U	K	S3	N5
Largescale stoneroller	<i>Campostoma oligolepis</i>	U	K	S3	N5
Western silvery minnow	<i>Hybognathus argyritis</i>	K	K	S1	N4
Mississippi silvery minnow	<i>Hybognathus nuchalis</i>	U	K	S3	N5
Plains minnow	<i>Hybognathus placitus</i>	C	S	S4	N4
Speckled chub	<i>Macrhybopsis aestivalis</i>	C	S	S3	N5
Sturgeon chub	<i>Macrhybopsis gelida</i>	R	K	SH	N3
Flathead chub	<i>Platygobio gracilis</i>	C	D	S3	N5
Sicklefin chub	<i>Macrhybopsis meeki</i>	R	K	S1?	N3
Gravel chub	<i>Erimytax x-punctatus</i>	U	K	S3	N4
Pallid shiner	<i>Hybopsis amnis</i>	R	D	S2	N4
Pugnose minnow	<i>Opsopoeodus emiliae</i>	U	K	S3	N5
Pugnose shiner	<i>Notropis anogenus</i>	R	K	S1	N3
Ghost shiner	<i>Notropis buchanani</i>	R	D	S2	N5
Blacknose shiner	<i>Notropis heterolepis</i>	R	K	S2	N4
Spottail shiner	<i>Notropis hudsonius</i>	C	S	S?	N5
Ozark minnow	<i>Notropis nubilus</i>	U	K	S3	N5
Weed shiner	<i>Notropis texanus</i>	R	D	S2	N5
Topeka shiner	<i>Notropis topeka</i>	R	D	S3	N3
Redfin shiner	<i>Lythrurus umbratilis</i>	U	D	S2	N5
Longnose dace	<i>Rhinichthys cataractae</i>	C	K	S3	N5
Pearl dace	<i>Margariscus margarita</i>	X		S1	N5
Southern redbelly dace	<i>Phoxinus erythrogaster</i>	C	D	S4	N5

Blue sucker	<i>Cycleptus elongatus</i>	C	D	S3	N3
Common Name	Scientific Name	Iowa Abundance	Iowa Trend	Iowa Status	National Status
Black buffalo	<i>Ictiobus niger</i>	U	K	S3	N5
Black redhorse	<i>Moxostoma duquesnei</i>	U	K	S3	N5
River redhorse	<i>Moxostoma carinatum</i>	R	K	S1	N4
Greater redhorse	<i>Moxostoma valenciennesi</i>	X	K	SX	N4
Spotted sucker	<i>Minytrema melanops</i>	C	K	S3	N5
Blue catfish	<i>Ictalurus furcatus</i>	U	S/D	S3	N5
Brown bullhead	<i>Ameiurus nebulosus</i>	R	D	S2	N5
Slender madtom	<i>Noturus exilis</i>	U	S	S3	N5
Tadpole madtom	<i>Noturus gyrinus</i>	U	D	S3	N5
Freckled madtom	<i>Noturus nocturnus</i>	R	K	S2	N5
Pirate perch	<i>Aphredoderus sayanus</i>	R	D	S3	N5
Trout perch	<i>Percopsis omiscomaycus</i>	U	D	S3	N5
Burbot	<i>Lota lota</i>	U	D	S3	N5
Banded killifish	<i>Fundulus diaphanus</i>	U	D	S2	N5
Blackstripe topminnow	<i>Fundulus notatus</i>	U	D	S3	N5
Mottled sculpin	<i>Cottus bairdi</i>	R	K	S2	N5
Slimy sculpin	<i>Cottus cognatus</i>	U	K	S3	N5
Slenderhead darter	<i>Percina phoxocephala</i>	U	K	S3	N5
Blackside darter	<i>Percina maculata</i>	C	D	S3	N5
River darter	<i>Percina shumardi</i>	C	S	S3	N5
Northern logperch	<i>Percina caprodes</i>	C	S	S3	N5
Crystal darter	<i>Crystallaria asprella</i>	R	K	S1	N3
Western sand darter	<i>Ammocrypta clara</i>	U	S	S2	N3
Banded darter	<i>Etheostoma zonale</i>	U	K	S3	N5
Bluntnose darter	<i>Etheostoma chlorosomum</i>	R	K	S1	N5
Mud darter	<i>Etheostoma asprigene</i>	U	S	S3	N4N5
Orangethroat darter	<i>Etheostoma spectabile</i>	U	K	S2	N5
Least darter	<i>Etheostoma microperca</i>	R	K	S1	N5

Table 3-9. Mussels of greatest conservation need.

Iowa Abundance: A = abundant, C = common, U = uncommon, R = rare, K = unknown, X = possibly extirpated.

Iowa Trend: K = unknown, I = increasing, S = stable, D = decreasing.

Iowa Listing: N = not listed, S = special concern, T = T, E = E, X = extirpated.

See Appendix 11 for NatureServe codes used in Iowa Status and National Status columns

Common Name	Scientific Name	Iowa Abundance	Iowa Trend	Iowa Listing	State Status	National Status
Elktoe	<i>Alasmidonta marginata</i>	U	D	N	S3	N4
Slippershell	<i>Alasmidonta viridis</i>	R	D	E	S1	N4
Flat floater	<i>Anodonta suborbiculata</i>	R	D	N	S1	N5
Cylinder	<i>Anodontoides ferussacianus</i>	R	D	T	S2	N5
Rock pocketbook	<i>Arcidens confragosus</i>	U	D	N	S3	N4
Spectacle case	<i>Cumberlandia monodonta</i>	R	D	E	S1	N2N3
Purple pimpleback	<i>Cyclonaias tuberculata</i>	R/X?	D	T	S1	N5
Butterfly	<i>Ellipsaria lineolata</i>	U	K	T	S2S3	N4
Spike	<i>Elliptio dilatata</i>	U	D	N	S2	N5
Ebonyshell	<i>Fusconaia ebena</i>	R	D	X	S1	N4N5
Ozark pigtoe	<i>Fusconaia ozarkensis</i>	E	X		SX	N3
Higgins' eye pearlymussel	<i>Lampsilis higginsii</i>	R	D	E	S1	N1
Yellow sandshell	<i>Lampsilis teres anodontoides</i>	R	D	E	S1	N5
Slough sandshell	<i>Lampsilis teres teres</i>	R	D	E	S1	N5
Creek heelsplitter	<i>Lasmigona compressa</i>	R	D	T	S1	N5
Fluted shell	<i>Lasmigona costata</i>	R	D	N	S2	N5
Pondmussel	<i>Ligumia subrostrata</i>	X	NA	X	SX?	N4N5
Hickorynut	<i>Obovaria olivaria</i>	U	D	N	S3	N4
Bullhead (Sheepnose)	<i>Plethobasus cyphus</i>	R	D	E	S1	N3
Round pigtoe	<i>Pleurobema sintoxia</i>	R	D	E	S2	N4
Monkeyface	<i>Quadrula metanerva</i>	U	D	N	S3	N4
Wartyback	<i>Quadrula nodulata</i>	U	D	N	S3	N4
Strange floater (Squawfoot)	<i>Strophitus undulatus</i>	R	D	T	S2	N5
Lilliput	<i>Toxolasma parvus</i>	R	D	N	S2	N5
Pistolgrip	<i>Tritogonia verrucosa</i>	R	D	E	S1	N4
Fawnsfoot	<i>Truncilla donaciformis</i>	R	D	N	S2	N5
Pondhorn	<i>Uniomerus tetralasmus</i>	R	K	N	S1	N4

Common Name	Scientific Name	Iowa Abundance	Iowa Trend	Iowa Listing	State Status	National Status
Paper pondshell	<i>Utterbackia imbecillis</i>	R	D	N	S2	N5
Ellipse	<i>Venustaconcha ellipsiformis</i>	R	D	T	S1	N3/4

Dragonflies and damselflies. The Dragonfly and Damselfly Working Group listed 106 species that occur in Iowa (Appendix 8) based primarily on Cruden and Gode (2000) with updates by working group member Ann Johnson. The full list consists of 103 species reported by Cruden and Gode (2000) as having been collected in Iowa since 1985, plus two accidentals and one new species that has been recently described from northeast Iowa. The list does not include 15 species listed by NatureServe as occurring in Iowa for which Cruden and Gode (2000) have not confirmed recent specimens.

The list of SGCN (Table 3-10) consists of 28 species with a state NatureServe status of S1 or S2. Eight species listed as S1 are known to have range distributions in Iowa that are not continuous with the species range in neighboring states. This may indicate that the overall range of the species is contracting. The complete list of Iowa dragonflies and damselflies and the list of species of greatest conservation need were both reviewed by the full Steering Committee.

Table 3-10. Dragonflies and damselflies of greatest conservation need.

Iowa Abundance: R = rare, ER = extremely rare.

See Appendix 11 for NatureServe codes used in Iowa Status and National Status columns

Common Name	Scientific Name	Iowa Abundance	Iowa Status	National Status
Rapids Clubtail	<i>Gomphus quadricolor</i>	ER	S1	N3N4
Spangled Skimmer	<i>Libellula cyanea</i>	ER	S1	N5
Slaty Skimmer	<i>Libellula incesta</i>	ER	S1	N5
Stygian Shadowdragon	<i>Neurocordulia yamaskanensis</i>	ER	S1	N5
Rusty Snaketail	<i>Ophiogomphus rupinsulensis</i>	ER	S1	N5
Sand Snaketail	<i>Ophiogomphus sp.</i>	ER	S1	not recorded
Mocha Emerald	<i>Somatochlora linearis</i>	ER	S1	N5
Brimstone Clubtail	<i>Stylurus intricatus</i>	ER	S1	N4
Blue-faced Meadowhawk	<i>Sympetrum ambiguum</i>	ER	S1	N5
Carolina Saddlebags	<i>Tramea carolina</i>	ER	S1	N5
Emma's Dancer	<i>Argia emma</i>	ER	S1	N5
Alkali Bluet	<i>Enallagma clausum</i>	ER	S1	N5
Elegant Spreadwing	<i>Lestes inaequalis</i>	ER	S1	N5
Sulphur-tipped Clubtail	<i>Gomphus militaris</i>	ER	S1S2	N5
Canada Darner	<i>Aeshna canadensis</i>	R	S2	N5
Variable Darner	<i>Aeshna interrupta</i>	R	S2	N5
Blue-eyed Darner	<i>Aeshna multicolor</i>	R	S2	N5
Green-striped Darner	<i>Aeshna verticalis</i>	R	S2	N5
Four-spotted Skimmer	<i>Libellula quadrimaculata</i>	R	S2	N5
Royal River Cruiser	<i>Macromia taeniolata</i>	R	S2	N5
Cyrano Darner	<i>Nasiaeschno pentacantha</i>	ER	S2	N5
Smoky Shadowdragon	<i>Neurocordulia molesta</i>	R	S2	N4
Paiute Dancer	<i>Argia alberta</i>	R	S2	N4
Prairie Bluet	<i>Coenagrion angulatum</i>	R	S2	N3?
Boreal Bluet	<i>Enallagma boreale</i>	R	S2	N5
Vesper Bluet	<i>Enallagma vesperum</i>	R	S2	N5
Spotted Spreadwing	<i>Lestes congener</i>	R	S2	N5
Sweetflag Spreadwing	<i>Lestes forcipatus</i>	R	S2	N5

SGCN in each Taxonomic Class

Table 3-11 compares the total number of species considered in each taxonomic class with the number of SGCN in each class. Birds and fish had the greatest number of species on the list, but each taxonomic class had some representatives. The classes with the highest percentage of species listed are those utilizing aquatic or semi-aquatic habitats – fish, mussels and dragonflies and damselflies.

Nearly all SGCN are nongame wildlife whose status is imperiled by lack or degradation of habitat. A few game animals with declining habitats or with declining trends nationally are listed e, g, ruffed grouse, woodcock, white-tailed jackrabbit, and bobwhite quail.

Nearly one third of all the wildlife considered by this Plan are considered to be in need of conservation to protect them from declining further into Threatened or Endangered status.

Table 3-11. Percent of Iowa species in each taxonomic class listed as a SGCN.

Group	Total Iowa Species Considered	Number with Greatest Need	Percent of Group Total
Fish	153	68	44
Breeding Birds	206	67	33
Butterflies	119	30	25
Mussels	55	29	53
Migratory Birds	199	18	9
Mammals	82	18	22
Land Snails ¹	8	8	100
Amphibians and Reptiles	71	31	44
Dragonflies and Damselflies	106	28	26
TOTALS	999	297	30

¹ Only Iowa T and E species are listed.