



US008105808B2

(12) **United States Patent**
Lorah et al.

(10) **Patent No.:** **US 8,105,808 B2**
(45) **Date of Patent:** **Jan. 31, 2012**

(54) **ANAEROBIC MICROBIAL COMPOSITION AND METHODS OF USING SAME**

(75) Inventors: **Michelle M. Lorah**, Finksburg, MD (US); **Elizabeth A. Jones**, Manassas, VA (US); **Mary A. Voytek**, Arlington, VA (US)

(73) Assignee: **The United States of America as represented by the Secretary of the Interior**, Washington, DC (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 582 days.

(21) Appl. No.: **12/131,666**

(22) Filed: **Jun. 2, 2008**

(65) **Prior Publication Data**

US 2008/0318303 A1 Dec. 25, 2008

Related U.S. Application Data

(60) Provisional application No. 60/936,017, filed on Jun. 13, 2007.

(51) **Int. Cl.**

C12P 7/00 (2006.01)
C12N 1/00 (2006.01)
C12N 1/12 (2006.01)

(52) **U.S. Cl.** **435/132; 435/243; 435/252.1**

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

6,503,746 B1 1/2003 Daane et al.
6,783,678 B2* 8/2004 Sorenson 210/610
6,946,248 B2 9/2005 Sowers et al.
2002/0015991 A1 2/2002 Brennan et al.
2006/0194303 A1 8/2006 DeWitt et al.
2006/0223160 A1 10/2006 Vanzin

OTHER PUBLICATIONS

Jones, Elizabeth J. P., et al., "Characterization of a Microbial Consortium Capable of Rapid and Simultaneous Dechlorination of

1,1,2,2-Tetrachloroethane and Chlorinated Ethane and Ethene Intermediates," *Bioremediation Journal*, 2006, p. 153-168, vol. 10.

Jones, M.M., et al., "Biostimulation and Bioaugmentation to Enhance Degradation of Chlorinated Solvents in Wetland Sediments," *In Situ and On-Site Bioremediation*, 2003.

Tiedje, James M., et al., "Aerobic and Anaerobic Transformation of cis-Dichloroethene (cis-DCE) and Vinyl Chloride (VC): Steps for Reliable Remediation," Final Technical Report CU-1167, Dec. 2003. GeoSyntec Consultants, "Bioaugmentation for Remediation of Chlorinated Solvents: Technology Development, Status, and Research Needs," Oct. 2005.

Washington Savannah River Company, "Enhancements to Natural Attenuation: Selected Case Studies," May 2007, WSRC-STI-2007-00250.

Jones, Elizabeth, J. P., et al., "Development and Composition of a Mixed Culture for Bioremediation of Chlorinated Ethenes and Ethanes," *In Situ and On-Site Bioremediation*, Jun. 2005.

Lorah, Michelle, et al., "Physiological Characterization of a Broad Spectrum Reductively Dechlorinating Consortium," *In Situ and On-Site Bioremediation*, Jun. 2005.

Lorah, Michelle, et al., "Electron Donor Preference of a Reductive Dechlorinating Consortium," *In Situ and On-Site Bioremediation*, Jun. 2005.

Majcher, Emily H., "Design and Installation of an Innovative Enhanced Bioremediation Pilot Test in a Tidal Wetland," *In Situ and On-Site Bioremediation*, Jun. 2005.

GeoSyntec Consultants, "Bioaugmentation for Remediation of Chlorinated Solvents: Technology Development, Status, and Research Needs," Oct. 2005.

Washington Savannah River Company, "Enhancements to Natural Attenuation: Selected Case Studies," May 2007, WSRC-STI-2007-00250.

* cited by examiner

Primary Examiner — Yong Pak

(74) *Attorney, Agent, or Firm* — Elizabeth Arwine

(57) **ABSTRACT**

A microbial composition for concurrent dechlorination of a mixture of chlorinated ethanes and chlorinated ethenes includes a isolated consortium of bioremediative microorganisms comprising strains of microorganism comprising *Clostridium*, *Acetobacterium*, *Dehalobacter*, *Bacteroides*, and *Proteobacteria*. The composition may also include *Methanomicrobia*.

9 Claims, 9 Drawing Sheets




FIG. 1




FIG. 2A




FIG. 2B




FIG. 3




FIG. 4A

FIG. 4B






FIG. 5A


FIG. 5B



FIGS. 6A-6B



FIGS. 7A-7D



FIGS. 8A-8B

1

ANAEROBIC MICROBIAL COMPOSITION AND METHODS OF USING SAME

This application claims priority of provisional patent application Ser. No. 60/936,017 filed on Jun. 13, 2007, the entirety of which is incorporated herein by reference.

I. FIELD OF THE INVENTION

The present invention relates to an anaerobic microbial composition and to methods of using the microbial composition for effectively dechlorinating at least one of chlorinated ethanes, chlorinated ethenes, chlorinated methanes, or mixtures thereof.

II. BACKGROUND OF THE INVENTION

Bioaugmentation (site inoculation with a microbial culture) is a proven approach for stimulating complete dechlorination of sites contaminated with chlorinated ethenes. However, cultures have not been available for the large-scale treatment of chlorinated ethane contamination. Of additional concern, chlorinated ethanes can inhibit the degradation of chlorinated ethenes. Thus, cultures are needed for bioremediation of sites with mixtures of these contaminants.

Contamination of groundwater with chlorinated ethenes and ethanes is a serious problem due to widespread and historic commercial, industrial, and military use, relative resistance to degradation, and associated health hazards. Under anaerobic conditions, chlorinated ethenes and ethanes can be partially reduced to less chlorinated compounds or completely degraded to nonchlorinated end products depending on the physiological capability of an indigenous microbial community.

Bacterial isolates capable of reducing 1,2-dichloroethane (DCA) and 1,1,1-trichloroethane have been identified. One isolate has been shown to reduce 1,1,2,2-tetrachloroethane (TeCA) to cis 1,2-dichloroethene (cisDCE). Recent research on a mixed culture demonstrated growth of *Dehalobacter* sp. with the reduction of 1,1,2-trichloroethane (TCA) to vinyl chloride (VC). See Jones et al., *Characterization of a Microbial Consortium Capable of Rapid and Simultaneous Dechlorination of 1,1,2,2-Tetrachloroethane and Chlorinated Ethane and Ethene Intermediates*, *Bioremediation Journal*, 10:153-168 (2006), the disclosure of which is incorporated herein by reference in its entirety.

III. SUMMARY OF INVENTION

According to an aspect of the invention, a composition is provided for concurrent dechlorination of a mixture of chlorinated ethanes and chlorinated ethenes. The composition includes an isolated bioremediative consortium comprising strains of microorganism comprising *Clostridium*, *Acetobacter*, *Dehalobacter*, *Bacteroides*, and *Proteobacteria*.

According to another aspect of the invention, a composition is provided for concurrent dechlorination of a mixture of chlorinated ethanes and chlorinated ethenes. The composition includes a non-naturally occurring consortium of dechlorinatingly effective microbial species. The consortium of effective dechlorinatingly microbial species comprises at least one 16S rDNA nucleic acid sequence that has more than 95% identity to a nucleic acid sequence consisting of SEQ ID NO 1, a nucleic acid sequence consisting of SEQ ID NO 2, or a nucleic acid sequence consisting of SEQ ID NO 3.

According to another aspect of the invention, a method for dechlorinating chlorinated waste is provided including con-

2

tacting at least one of chlorinated ethanes, chlorinate ethenes, or chlorinated methanes with an isolated bioremediative consortium comprising strains of microorganism comprising Clostridiales, *Cytophaga-flavobacterium-bacterioides*, *Proteobacteria*, and *Methanomicrobia*; and anaerobically dechlorinating the at least one of chlorinated ethanes, chlorinate ethenes or chlorinated methanes.

According to another aspect of the invention, a method of producing a microbial consortium comprises culturing microbes of a sediment sample obtained from a site contaminated with a mixture of chlorinated ethanes and chlorinated ethenes in an anaerobic medium with at least one chlorinated ethane and an electron donor.

IV. BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 illustrates possible pathways of anaerobic 1,1,2,2-tetrachloroethane (TeCA) dechlorination. Compounds for which the EPA reports an increased risk of cancer are indicated with an asterisk.

FIGS. 2A-2B illustrate dechlorination in stock microbial cultures according to the present invention amended with lactate and (A) TeCA only or lactate and (B) TeCA, TCA and cisDCE.

FIG. 3 illustrates degradation of ethene and production of ethane in a microbial culture according to the present invention after depletion of chlorinated compounds.

FIGS. 4A-4B illustrate cisDCE (A) and TCA (B) degradation by stock microbial cultures according to the present invention grown with (1) TeCA only and (2) a mixture of TeCA, TCA, and cisDCE, and (3) no culture added.

FIGS. 5A-5B illustrate production and degradation of chlorinated intermediates of (A) TCA and (B) cisDCE in a microbial culture according to the present invention.

FIGS. 6A-6B illustrate dechlorination of added compounds and accumulation of intermediates in cultures transferred to fresh medium and amended with lactate and either (A) TeCA only or (B) TeCA, TCA, and cisDCE.

FIGS. 7A-7D illustrate a comparison of TRFLP profiles for APG sediments WB23 (A) and WB30 (B), and for a microbial composition according to the present invention after 1 year in culture with TeCA (C) or a mixture of chlorinated compounds (D).

FIGS. 8A-8B illustrate frequency of phylogenetic types in a microbial composition rDNA clone library (A) and mcr clone library (B) according to the present invention.

V. DETAILED DESCRIPTION OF INVENTION

The present invention is directed to an anaerobic microbial composition or consortium comprising bioremediative organisms. The invention is also directed to methods of using the microbial composition for the effective dechlorination of at least one of chlorinated ethanes, chlorinated ethenes, chlorinated methanes, or mixtures thereof. The consortium of the present invention may be employed for bioremediation to anaerobically biodegrade chlorinated waste, for example, contaminated groundwater or contaminated soil from landfill sites, river beds, lakes, wetlands, and the like.

FIG. 1 illustrates that the TeCA degradation pathway is primarily biotic and includes both hydrogenolysis to less chlorinated ethanes and dichloroelimination to less chlorinated ethenes. Abiotic production includes the production of trichloroethene (TCE) from dehydrochlorination of TeCA. Several intermediates of TeCA dechlorination are possible carcinogens and are listed as contaminants of concern by the U.S. Environmental Protection Agency. VC is a known

human carcinogen that often accumulates at sites where dechlorination is slow or incomplete.

Natural attenuation of TeCA has been documented at Aberdeen Proving Ground (APG) in Maryland, where contaminated groundwater discharges through anoxic wetland sediments at West Branch Canal Creek. Chloroethane (CA), ethene and ethane have not been observed in the sediment. According to the present invention, a microbial consortium, West Branch Consortium (WBC-2), was derived from organic-rich sediments collected in the wetland of West Branch Canal Creek at APG. As disclosed below, WBC-2 can dechlorinate chlorinated ethanes and chlorinated ethenes, both individually and concurrently.

According to an embodiment of the present invention, a microbial composition, WBC-2, for concurrent dechlorination of a mixture of chlorinated ethanes and chlorinated ethenes, comprises an isolated bioremediative consortium comprising strains of microorganism comprising *Clostridium*, *Acetobacter*, *Dehalobacter*, *Bacteroides*, and *Proteobacteria*. The microbial composition may also comprise Methanomicrobia, for example, at least one of *Methanosarcina* or *Methanosaeta*. The consortium may comprise *Dehalococcoides*.

In embodiments, the microbial composition may comprise a non-naturally occurring consortium of dechlorinatingly effective microbial species comprises at least one 16S rDNA nucleic acid sequence that has more than 95% identity to a nucleic acid sequence consisting of SEQ ID NO 1, a nucleic acid sequence consisting of SEQ ID NO 2, or a nucleic acid sequence consisting of SEQ ID NO 3. In embodiments, the consortium may comprise at least one 16S rDNA nucleic acid sequence that has more than 95% identity to a nucleic acid sequence consisting of SEQ ID NO 4 or SEQ ID NO 5. In embodiments, the consortium may comprise at least one 16S rDNA nucleic acid sequence that has more than 95% identity to a nucleic acid sequence consisting of SEQ ID NO 6 or SEQ ID NO 7.

In embodiments, the microbial composition may comprise at least one methyl coenzyme-M reductase nucleic acid sequence that has more than 95% identity to a nucleic acid sequence consisting of SEQ ID NO 8, SEQ ID NO 9, SEQ ID NO 10, SEQ ID NO 11, SEQ ID NO 12, SEQ ID NO 13, SEQ ID NO 14, SEQ ID NO 15, SEQ ID NO 16, SEQ ID NO 17, SEQ ID NO 18, SEQ ID NO 19, or SEQ ID NO 20.

According to a method of the present invention, a mixture of chlorinated ethanes and chlorinated ethenes may be dechlorinated by contacting the mixture with a microbial composition of the present invention and concurrently anaerobically dechlorinating the mixture of chlorinated ethanes and chlorinated ethenes. According to an embodiment of the invention, the chlorinated ethanes may comprise at least one of 1,1,2,2-tetrachloroethane; 1,1,2-trichloroethane; 1,2-dichloroethane, or chloroethane. The chlorinated ethenes may comprise at least one of cis 1,2-dichloroethene; trans 1,2-dichloroethene; vinyl chloride; or tetrachloroethene. The mixture may also contain chlorinated methane, for example, carbon tetrachloride or chloroform.

According to a method of the present invention, chlorinated waste may be dechlorinated by contacting at least one of chlorinated ethanes, chlorinate ethenes or chlorinated methanes with an isolated bioremediative consortium comprising strains of microorganism comprising *Clostridiales*, *Cytophaga-flavobacterium-bacterioides*, *Proteobacteria*, and *Methanomicrobia*; and anaerobically dechlorinating the at least one of chlorinated ethanes, chlorinate ethenes, or chlorinated methanes. The chlorinated waste may comprise contaminated soil or contaminated water.

The microbial consortium according to the present invention may be produced by culturing microbes of a sediment sample obtained from a site contaminated with a mixture of chlorinated ethanes and chlorinated ethenes in an anaerobic medium with at least one chlorinated ethane and an electron donor. In embodiments, the electron donor may be selected from the group consisting of lactate, pyruvate, benzoate, and ethanol.

Example

Properites and Composition of WBC-2 Anaerobic Microbial Composition

A. Dechlorination Properties of WBC-2 Microbial Composition

WBC-2 was cultivated in batches amended with either lactate and TeCA or lactate and a mixture of TeCA, TCA, and cisDCE for 18 months.

In the TeCA amended culture, WBC-2 in a 2 L batch culture completely dechlorinated TeCA (measured at 240 μM) within 2 days, as shown in FIG. 2A. The pathway of TeCA degradation could not be discerned by monitoring this culture because very little intermediate accumulation occurred. Less than 0.5 μM (0.2% of the added TeCA) accumulated as VC, and this VC was degraded by day 2. The end products of dechlorination were ethene and ethane.

In a 2 L batch culture maintained with the chlorinated mixture (TeCA, TCA, and cisDCE), WBC-2 rapidly, simultaneously, and completely reduced all three chlorinated compounds to the nonchlorinated end-products ethene and ethane, as shown in FIG. 2B. Small amounts of transDCE and VC were observed as transient intermediates.

After dechlorination was complete, the fate of ethene was monitored in the cultures in order to determine if ethene reduction could account for the production of ethane. As shown in FIG. 3, ethene was degraded (6 $\mu\text{M day}^{-1}$) with the production of ethane. After ethene was depleted, the ethane concentration also decreased, at a rate of 0.5 $\mu\text{M day}^{-1}$. The lack of stoichiometric accumulation of ethane suggests that ethane and ethene degradation can co-occur.

Pathways of intermediate degradation were determined by incubating sub samples of WBC-2 with TCA or cisDCE. WBC-2 cultures that had been maintained with (1) TeCA and (2) the chlorinated mixture (TeCA, TCA, cisDCE) were incubated to deplete chlorinated compounds and then compared with respect to their capabilities to degrade cisDCE (FIG. 4A) and TCA (FIG. 4B). The two cultures were very similar in their abilities to degrade the two TeCA intermediates.

Degradation products of intermediate dechlorination are shown for TeCA-maintained WBC-2 (FIGS. 5A-5B). Little intermediate accumulation was observed in TCA-amended treatments (FIG. 5A). The rate of TCA degradation was 36 $\mu\text{M day}^{-1}$. The peak VC and DCA concentrations measured were 1.7% and 0.3% of the TCA added, respectively, and both intermediates were rapidly degraded. Chloroethane was not detected. As shown in FIG. 5B, cisDCE reduction was accompanied by the production of VC (peak accumulation, 24% of added cisDCE), ethene and ethane. The rate of cisDCE degradation in cultures amended with cisDCE was 54 $\mu\text{M day}^{-1}$.

When WBC-2 cultures were diluted (1:9) into fresh medium, dechlorination was initially slow enough to allow the observation of intermediates. As shown in FIG. 6A, diluted cultures amended with lactate and TeCA (as shown by arrows) accumulated measurable TCA (0.01 μM) at one time point, and transDCE (6 μM) and VC (0.25 μM) were also both produced and degraded.

5

As shown in FIG. 6B, diluted cultures amended with TeCA, TCA, and cisDCE (as shown by arrows) had a transient accumulation of VC (6% of the added chlorinated compounds) and transDCE (as much as 4% of added TeCA), and also the abiotic product, TCE (3% of the TeCA parent added). Within 2 to 4 weeks, dechlorination rates increased to a level comparable to that observed in undiluted cultures, with parent compounds degraded in as little as one day and little transient accumulation of intermediates.

After microbial enrichment and 1 year in culture, the rates of WBC-2 dechlorination for TeCA, TCA, and cisDCE were 100, 36, and 54 $\mu\text{M day}^{-1}$, respectively. Almost no intermediates were detected in WBC-2 cultures amended with TeCA or a mixture of TeCA, TCA, and cisDCE. Although intermediates often did not accumulate, the ability of WBC-2 to degrade all known intermediates in the TeCA pathway was demonstrated in treatments with individual compounds (e.g., TCA and cisDCE) and in mixtures (e.g., TeCA, TCA, cisDCE). Ethene and ethane are major products of TeCA degradation.

The unique contribution of WBC-2 for contaminant treatment is the ability of the microbial composition to handle a variety of compounds (both chlorinated ethenes and ethanes) without noticeable inhibition. WBC-2 also has the ability to degrade chlorinated ethenes and ethanes simultaneously. WBC-2 is also able to reduce tetrachloroethene (PCE) in the presence of TeCA.

WBC-2 also has other properties regarding sensitivity to pH, oxygen, and chlorinated methanes.

Batch microcosms were conducted under anaerobic conditions in culture medium with neutral pH and with pH adjusted from acidic (pH 4, 5, and 6) to alkaline (pH 8 and 9). The ability of the WBC-2 consortium to fully dechlorinate chlorinated ethanes and ethenes was evaluated for each pH-adjusted treatment. The WBC-2 consortium was intolerant of acidic conditions of pH 5 and lower, resulting in a loss of dechlorinating ability. The consortium was tolerant of alkaline pH with no apparent loss of activity.

Reductive dechlorinating organisms commonly have high negative sensitivity to oxygen exposure over short timeframes, which complicates their field application. To evaluate oxygen sensitivity of WBC-2, an aliquot was removed from an anaerobic culture vessel and poured into smaller containers on the bench top where a series of oxygen exposures were applied to the culture by bubbling ambient air through the culture at a rate of approximately 100 milliliters per minute. Following time exposures of 1, 5, 20, and 60 minutes, each treatment was purged with anaerobic gas, sealed, and

6

amended with 1,1,2,2-tetrachloroethane, 1,1,2-trichloroethane, and cis-1,2-dichloroethene, and an electron donor. Treatments were monitored for dechlorinating for 11 days or until complete dechlorination was observed. Dechlorinating activity was observed to be comparable to the culture with no exposure to oxygen up to an exposure time of 20 minutes. After 60 minutes of exposure, dechlorination activity was partially inhibited but eventually continued to completion during incubation under strict anaerobic conditions.

Chlorinated methanes tend to inhibit the activity of a wide range of organisms. To test the sensitivity of WBC-2 to carbon tetrachloride (CT), a series of batch experiments in culture medium were conducted along with flow-through column experiments with a bioaugmented, wetland-like, organic matrix. Although toxicity effects from CT addition were observed with WBC-2 in liquid culture at a concentration of 3 mg/L, WBC-2 in the columns could maintain degradation of CT and chloroform (CF) and of the chlorinated ethanes and ethenes at CT and CF concentrations of 10 and 20 mg/L, respectively.

B. Composition of the WBC-2 Microbial Composition

As shown in FIGS. 7A-7D, the bacterial community in the source APG sediments (WB23, FIG. 7A; WB30, FIG. 7B) shifted after a year under culture conditions with lactate as the electron donor and TeCA (FIG. 7C) or a mixture of TeCA, TCA, and cisDCE (FIG. 7D) as electron acceptors.

Both WBC-2 cultures (cultured with TeCA as shown in FIG. 7C or TeCA, TCA, and cisDCE as shown in FIG. 7D) exhibited TRFLP profiles that were different from that of the source sediments and overall represented a different community than was present in the starting materials. This change reflects the selection pressures exerted on the community and individual members by the chlorinated compounds, such that the remaining peaks represent members tolerant of the chlorinated compounds and favored by the culturing conditions and perhaps directly or indirectly involved in the degradation process.

The numerically dominant phylogenetic types in WBC-2 were identified by cloning and sequencing 16S rDNA and *mcrA* genes from a culture grown with a mixture of TeCA, TCA, and cisDCE. The frequency of phylotype occurrence in 16S rDNA clone libraries and *mcrA* clone libraries, and phylogenetic placement was determined using a BLAST search for related sequences.

Although most of the WBC-2 clones were not related to dechlorinating bacteria that have been studied in isolation, many were related to bacterial clones that have been observed at other dechlorinating sites, as shown in Table 2:

TABLE 2

Frequency of WBC-2 Clones and Their Closest BLAST Matches to a Dechlorinating Isolate or to Clone Sequences from a Dechlorinating Environment				
Phylotype frequency	Phylogenetic placement	Clone id	BLAST hits ¹ (% similar)	Source
48/133	Clostridium	acc #DQ907198	*AY667266 (99%)	TCB-dechlorinating community
35/133	Acetobacteria	acc #DQ907202	*AY185312 (97%) AY185315 (97%) AY185311 (96%)	1,2-Dichloropropane-dechlorinating enrichment
2/133	Dehalobacter	acc #DQ907207	*AF422637 (95%) AY754830 (93%) DQ663785 (93%)	TCB-reducing community PCB-dechlorinating culture 111-trichloroethane degrading mixed culture
19/133	Bacteroides	acc #DQ907199	*DQ080146 (95%) AY553955 (95%)	2,3,4,5-Tetrachlorobiphenyl culture containing Dehalococcoides PCB contaminated harbor sediment

TABLE 2-continued

Frequency of WBC-2 Clones and Their Closest BLAST Matches to a Dechlorinating Isolate or to Clone Sequences from a Dechlorinating Environment				
Phylotype frequency	Phylogenetic placement	Clone id	BLAST hits ¹ (% similar)	Source
1/133	Geobacter	acc #DQ907201	AY780553 (95%)	Chlorinated ethene-dechlorinating enrichment
			AJ488070 (90%)	Chlorobenzene degrading consortium
			*AY217446 (97%)	TCE-dechlorinating community
			*AY217435 (97%)	TCE-dechlorinating community
			*AY780563 (98%)	Chlorinated ethene enrichment culture
			AF223382 (98%)	Isolate that dechlorinates trichloroacetic acid
			AF447133 (98%)	Population that dechlorinates saturated PCE
7/133	Pseudomonas	acc #DQ907203	AY914177 (98%)	Isolate that dechlorinates PCE
			AY667270 (97%)	TCE contaminated aquifer
			AY017341 (99%)	Chlorate-reducing isolate

¹Top BLAST hit.

As shown in FIG. 8A, the 16S rDNA library was dominated by Clostridiales (65%), including three phylotypes. The phylotype representing the greatest number of clones was a *Clostridium* sp. (SEQ ID NO 1), most closely related (99%) to an uncultured member of a TCE dechlorinating community. The second most prevalent phylotype was an *Acetobacterium* sp. (SEQ ID NO 2) most closely related (97%) to uncultured clones from a 1,2-dichloropropane-dechlorinating enrichment, and 97% and 96% related to the homoacetogens *Acetobacterium malicum* and *A. wieringae*, respectively. Less prevalent was a third phylotype, 95% related to an uncultured clone from a TCE-dechlorinating community and 93% related to *Dehalobacter restrictus* (SEQ ID NO 3), in the evaluated region between 46f and 519r.

There was more variability among sequences of the Bacteroidetes (Cytophaga-flavobacterium-bacteroides group), which accounted for 14% bacterial clones (SEQ ID NO 4 and SEQ ID NO 5). Many of these were related to uncultured clones from dechlorinating populations (Table 2).

Among the Proteobacteria, one clone (delta Proteobacteria) was most closely related (98%) to a *Geobacter* sp. (SEQ ID NO 6) from a chlorinated ethene enrichment culture, and 98% related to *G. lovleyi*, a recently described PCE-dechlorinating isolate. The gamma Proteobacteria (SEQ ID NO 7) were 99% related to the top 60 BLAST hits, including *Pseudomonas stutzeri* and *Ps. chloridismutans*, that is able to dechlorinate trichloroacetic acid.

For the other phylotypes observed among the bacterial clones (*Arcobacter* sp. and *Desulfobulbus* sp., 14% and 2% of the clones, respectively) and all of the *mcrA* clones, the BLAST database did not reveal relatedness to organisms from dechlorinating populations. *Arcobacter* sp. comprises part of the epsilon proteobacteria. *Desulfobulbus* sp., comprises part of the delta proteobacteria.

No *Dehalococcoides* clones were identified. However, *Dehalococcoides* numbers determined independently using qPCR and compared with the total number of cells by microscopic count indicated that about 1% of the total consortium population was comprised of *Dehalococcoides* spp. Microscopic examination confirmed that cocci were rare, and the consortium population was composed almost entirely of rod-shaped cells.

As shown in FIG. 8B, the WBC-2 *mcrA* clone library was comprised of members of the class Methanomicrobia, and included both acetate- and H₂-utilizing methanogens (acces-

sion numbers DQ907209 to DQ907221, i.e., SEQ ID NO 8, SEQ ID NO 9, SEQ ID NO 10, SEQ ID NO 11, SEQ ID NO 12, SEQ ID NO 13, SEQ ID NO 14, SEQ ID NO 15, SEQ ID NO 16, SEQ ID NO 17, SEQ ID NO 18, SEQ ID NO 19, and SEQ ID NO 20). Members of the Methanosarcinaceae family are capable of utilizing acetate in the production of methane. These include *Methanosarcina* spp., which may utilize acetate, H₂, methanol, or methyl amines, and *Methanosaeta* spp., which are obligate acetate utilizers. The other Methanomicrobia are related to methanogens that utilize H₂ and formate as electron donors. The presence of *Methanosaeta* spp. in the WBC-2 culture indicates that acetate is being produced. Although the cultures were methanogenic, methanogens comprised a very small part of the total microbial population of WBC-2. Total methanogens quantified using qPCR comprised 0.2% of the total WBC-2 microbial population.

C. Evidence for a Distinct Dechlorinating Population

The TeCA dechlorination pathway includes reactions that are a part of the more thoroughly studied PCE dechlorination pathway (right half of FIG. 1), and thus might be expected to support the growth of similar organisms. Indeed, the predominance of Clostridia and CFB in the WBC-2 population (Table 2) is also characteristic of TCE-dechlorinating communities, and organisms similar to the ethene-dechlorinating *Dehalococcoides* and *Dehalobacter* are observed in WBC-2.

However, the dechlorinating abilities of WBC-2, enriched in the presence of chlorinated ethanes, are different than cultures enriched with chlorinated ethenes, both with respect to the response to added electron donors and in the relative importance of *Dehalococcoides* within the microbial population. Based on limited studies, the electron donor needs for chlorinated ethane-enriched cultures appear to be different from those of cultures enriched for chlorinated ethene reduction.

Most bacterial isolates capable of reductive dechlorination of chlorinated ethenes, including *Dehalococcoides* and *Dehalobacter*, use H₂ as the preferred electron donor. The chlorinated ethane-enriched culture, WBC-2, was not stimulated to reduce cisDCE or TCA with H₂, H₂ plus acetate, or propionate added as the electron donor. The failure of H₂ to stimulate dechlorination suggests that the organisms involved are not the same as the chlorinated ethene-reducing organisms that have been studied in isolation. In addition, *Dehalococcoides* spp. comprise only a minor part (about 1%) of the

cell population in WBC-2. This, coupled with the presence of consortium members that appear to be closely related to clones of unknown function in other dechlorinating populations, suggests that organisms other than *Dehalococcoides* spp. may play a greater role in TeCA dechlorination. These observations suggest that exposure to chlorinated ethanes results in the selection of a different population of organisms for chlorinated ethene reduction. Cultures that were enriched using chlorinated ethenes (i.e., in the absence of chlorinated ethanes) were inhibited in the presence of chlorinated ethanes.

D. Microbial Roles Within WBC-2

The specific roles of WBC-2 consortium members have yet to be determined. Although tentative roles could be assigned to *Dehalococcoides* (cisDCE, transDCE, VC, and DCA reduction), *Dehalobacter* (TCA reduction to VC, DCA reduction to ethene), and *Acetobacterium* (DCA reduction to ethene) based on studies of related organisms, WBC-2 exhibited some capabilities, such as TeCA reduction to transDCE, and TCA reduction to DCA, for which no organisms have been implicated.

The observation of closely related phylotypes in WBC-2 and other dechlorinating communities, such as *Clostridium* sp., *Acetobacterium* sp., and CFB provides some evidence for the involvement of previously unrecognized bacteria in dechlorination processes. However, enrichment of organisms in a dechlorinating system provides only circumstantial evidence for direct involvement. In WBC-2, lactate fermentation, homoacetogenesis, methanogenesis, sulfur cycling, syntrophy, and chemoautotrophy could support organisms in the culture without deriving energy from dechlorination. The possible role of methanogens in dechlorination by WBC-2 is not known. Although the presence and activity of methanogens are often considered to be inhibitory to or at least contraindicative of dechlorination activity, some evidence suggests that methanogens may play an important role in dechlorination.

Example

Bioremediation with WBC-2

WBC-2 was able to degrade chlorinated ethanes and ethenes simultaneously with little VC accumulation. This capability makes the microbial consortium WBC-2 a potentially valuable tool for bioremediation of sites contaminated with mixtures of chlorinated ethenes and ethanes. In addition, the simultaneous reduction of all components of the TeCA degradation pathway can reduce the total treatment time and help prevent transport of hazardous compounds out of the treatment zone.

An innovative, enhanced bioremediation pilot test was designed and installed in a seep at the ground-water/surface-water interface within the tidal wetland along West Branch Canal Creek at Aberdeen Proving Ground, Maryland, to treat a mixture of dissolved chlorinated methanes, ethenes, and ethanes.

First-order degradation rates were estimated by circulating increasing concentrations of site contaminant mixtures through continuous, up-flow columns comprised of a commercially available compost, peat, and sand-matrix (with and without zero-valent iron).

Columns were bioaugmented with a mixed anaerobic consortium (WBC-2). The tolerable thickness of the bioreactive mat at this site was determined based on sediment strength, bearing capacity, and estimated settlement over time. Thus, the bioreactive mat was comprised of two reactive zones: (1)

a lower zone (nearest the sediment) designed to enhance abiotic degradation of chlorinated methanes, and (2) an upper zone designed to enhance biodegradation of the remaining chlorinated methanes, ethanes, and ethenes. The lower zone was comprised of zero-valent iron filings and an organic matrix consisting of commercially-available compost, peat, and sand. The upper zone was comprised of a compost, peat, and sand matrix and was bioaugmented with WBC-2.

Example

Methods and Materials

I. Development of WBC-2 Anaerobic Microbial Composition

Sediment was collected from two sites within the wetland at West Branch Canal Creek (WB23 and WB30) and prepared anaerobically. Sediments were collected, sieved, slurried with groundwater (1:1.5), and incubated (19° C.) with TeCA (7 μM) in 1-L serum bottles without headspace for 1 month. Most of the Fe(III) and sulfate (alternative electron acceptors) were depleted during this incubation period and methane was being produced. Aliquots (100 ml) of sediment slurries were then transferred to 120-ml serum bottles with a N₂/CO₂ (95:5) headspace and amended with a daughter compound, cisDCE or TCA, for 1 to 2 months. In all sediment slurry enrichments, the electron donors were derived only from organic matter in the sediment.

Sediment slurries (100 ml each of TCA-enriched WB23 and cisDCE-enriched WB30) were then transferred into an anaerobic culture medium as described below (1800 ml) with sulfide (50 μM) added as a reductant, and amended with target concentrations of TeCA (30 μM) or a mixture of TeCA (25 μM), TCA (50 μM), and cisDCE (50 μM). The electron donor for cultivation was selected in tests on the TeCA amended culture (see below) 3 weeks after inoculation from sediment slurry. Cultures were diluted over a 2-year period and contain about 0.1% sediment by volume.

II. Evaluation of Electron Donors

All electron donor tests were performed in duplicate on sub samples removed from TeCA-depleted stock culture. Because concentrations of intermediates often are low or undetectable during TeCA degradation by WBC-2, TCA and cisDCE were used as test compounds to ensure that the electron donor selected would support both chlorinated ethane and chlorinated ethene pathways. In addition, the ability of each electron donor to support the dechlorination of VC was evaluated.

Aliquots of WBC-2 culture (10 ml) were transferred anaerobically to 28-ml pressure tubes filled with N₂/CO₂ (80:20). WBC-2 was evaluated for dechlorination of test compounds in the following electron donor treatments: propionate (10 mM); succinate (3 mM); lactate (3 mM); pyruvate (3 mM); benzoate (3 mM); formate (10 mM); acetate (10 mM); H₂ (20 kPa overpressure, added three times during the incubation) with or without acetate (1 mM) added as a carbon source; whey (5 g/L); no electron donor added.

The electron donors supplied electron equivalents (assuming complete oxidation) equal to about 100 times that required for the reduction of the chlorinated compounds, cisDCE or TCA, which were added from aqueous emulsions (for a final concentration of approximately 1 and 0.75 mM, respectively) and monitored for VC and DCA production. The ability to dechlorinate VC was tested by adding VC (4.2 μM) from a gaseous standard (Matheson, Twinsburg, Ohio) in a separate treatment. All treatments were incubated at 19° C.

and monitored by sampling the headspace for analysis with a gas chromatograph (GC) with a flame ionization detector (FID).

As shown in Table 1 below, the electron donor for WBC-2 cultivation was selected by comparing dechlorination in electron donor treatments with dechlorination in a control with no added electron donor. Controls with no added electron donor exhibited decreases in added TCA and cisDCE characteristic of adsorption, with an initial decrease of 24% and 10%, respectively, followed by no further decrease. Less than 1% of the added TCA or cisDCE in the controls was reduced to VC and no DCA was produced. H₂ did not stimulate reduction of TCA or cisDCE above that observed in the controls, whether or not acetate was added as a carbon source.

The most complete dechlorination was obtained in treatments with lactate and pyruvate. The pathway of TCA reduction (production of VC versus DCA from TCA) varied among electron donor treatments. Cultures with lactate produced more DCA relative to VC than did cultures with pyruvate. No DCA was produced in treatments with propionate, acetate, benzoate, or whey.

TABLE 1

WBC-2 Reduction of cDCE, TCA and VC			
Electron donor	Degradation relative to no electron donor		
	cDCE	TCA	VC
Succinate	+	+/-	-
Lactate	++	+++	+++
Pyruvate	+	++	+++
Benzoate	-	+	ND
Propionate	+	-	-
Formate	-	-	+
Acetate	-	-	-
H ₂	-	-	+++
H ₂ , with 1 mM acetate	-	-	+++
Whey	-	-	ND

+++ , complete dechlorination (TCA [0.75 mM], DCE [1 mM] after 27 days, or VC [4.2 μM] after 3 days).

++ , dechlorination at least 50% complete.

+ , greater than no electron donor control.

+/- , one duplicate greater than control.

- , dechlorination not greater than no electron donor control.

ND, treatment not done.

III. Culture Medium and Maintenance

The anaerobic medium included (g/L deionized water): NaHCO₃ (2.5), NH₄Cl (0.5), NaPO₄ (0.5), KCl (0.1), 10 ml vitamin solution, and 10 ml trace mineral solution, with a gas phase of N₂ and CO₂ (80:20). The trace mineral solution contained (g/L): Nitritotriacetic acid (1.5), MgSO₄·7H₂O (3.0), MnSO₄·H₂O (0.5), NaCl (1.0), FeSO₄·7H₂O (0.1), CaCl₂·2H₂O (0.1), COCl₂·6H₂O (0.1), ZnCl₂ (0.13), CuSO₄·5H₂O (0.01), AlK(SO₄)₂·12H₂O (0.01), H₃BO₃ (0.01), Na₂MoO₄ (0.025), NiCl₂·6H₂O (0.024), Na₂WO₄·2H₂O (0.025).

Early in development, some batches were starved for periods as long as several months, but recovered activity when feeding was resumed. Once established, cultures were maintained either with lactate (1 mM) and TeCA (50 μM) added from aqueous stocks once or twice weekly, or lactate (1.5 mM) and TeCA (25 μM), TCA (50 μM), and cisDCE (50 μM). The ratio of electron equivalents for donors to acceptors was 30 and 17, respectively, for stocks maintained with TeCA and the chlorinated mixture. The chlorinated stocks were prepared by adding purified standards to sterile anaerobic deionized water. Chlorinated stock bottles were vigorously shaken

prior to each amendment in order to emulsify any undissolved compound. The cisDCE contained approximately 1% transDCE.

WBC-2 was maintained in 2 L batches of medium from which culture volumes were removed for study. After 1 year, all cultures were restored to full 2 L volume by adding fresh medium. Under contract by the U.S. Army, samples of WBC-2 were given to SIREM Laboratories (Guelph, Ontario, Canada) for propagation of the culture to the large volumes required for field bioremediation tests conducted by the U.S. Geological Survey, using lactate and the chlorinated mixture described above (GeoSyntec Consultants Inc., 2004). WBC-2 microbial composition (clone analysis) was assessed in a culture sample obtained from SIREM Laboratories that had been scaled up by transferring two times into fresh medium.

IV. Gas Analysis

Culture headspace was sampled using a gas tight syringe and injected into one or both of the following GC-FID systems. For rapid analysis of TeCA, TCA, DCA, TCE, cisDCE, transDCE, and VC, a Hewlett-Packard model 5890 series 11 with isothermal separation at 100° C. on a VOCOL (Supelco, Bellefonte, Pa.) capillary column (30 m×0.53 mm) was used. For separation of methane, ethene, and ethane, and for analysis of VC, transDCE, and cisDCE when interfering peaks were present, a Shimadzu model GC-17A with separation on a RtQ-Plot (Restek, Bellefonte, Pa.) column (30 m×0.32 mm) using a temperature program of 100° C. for 5 min, ramping to 200° C. at 20° C./min was used.

Aqueous standards of chlorinated compounds were prepared from highly purified neat calibration standards. Standards of chlorinated compounds were prepared by adding 10 μl of neat solution to 100 ml water, and preparing aqueous dilutions for headspace analysis in bottles sealed with Teflon coated stoppers. Dimensionless Henry's law constants (DHLCs), were used to calculate expected headspace concentrations from known liquid concentrations. Methane, ethene, and ethane standards were purchased as gas standards (Scott Specialty Gas). Concentrations of chlorinated compounds and non-chlorinated end products in samples of WBC-2 culture headspace were converted to dissolved values using DHLCs and total concentrations (per volume medium) were calculated. DHLCs have been measured empirically by many researchers and vary widely. The chosen DHLCs fall in the midrange of published values. Nonetheless, the DHLCs are the greatest source of possible error in the concentrations reported here, and may exceed 10%. Errors between repeat injections are about 2%.

The dimensionless Henry's law constants (DHLC) applied were 0.019 for TeCA, 0.556 for TCA, 0.1821 for DCA, 0.3056 for TCE, 0.1255 for cisDCE, 0.3056 for transDCE, 0.9087 for VC, 7.96 for ethene, 19.88 for ethane, and 28.5 for methane. All detection limits were less than 0.01 μM.

V. Monitoring WBC-2 Dechlorination

A simple method was used for monitoring dechlorination, based on partitioning in the headspace. However, the stoichiometry of TeCA to its degradation products is difficult to determine using this method. Both adding and measuring the initial concentration of TeCA involved potentially large errors. Dilution of the stock solution cannot be relied upon to determine the amount of TeCA added, because TeCA in the stock solution was not completely dissolved and possibly not completely homogenized by shaking. TeCA measurements were subject to two known sources of error. First, when calculating total concentrations from headspace values, errors are magnified for compounds with a very low DHLC. The DHLC for TeCA (0.019) was an order of magnitude lower than any other compound measured and therefore sub-

ject to the greatest error. Conversely, compounds such as methane, ethene and ethane are largely partitioned to the headspace and subject to the least error. Second, because there is some sediment as well as cell mass in the cultures, distribution of TeCA between the liquid and gas phases may be complicated by adsorption, resulting in a discontinuity between TeCA uptake and the appearance of end product. All of these errors may have contributed to differences between target TeCA additions and measured concentrations (e.g., target addition of 60 μM versus measured concentration, 240 μM , that is reported in FIG. 2), as well as the error in stoichiometry between different bottles (i.e., 2 ± 1).

VI. Cloning, RFLP Screening and Sequencing, TRFLP

DNA was extracted using the Bio-101 Fast DNA Spin Kit for Soil (MP Biomedicals, Irvine, Calif.) following manufacturers instructions, except that product recovery was maximized at each step. Bacterial and methanogen DNA were amplified using the polymerase chain reaction (PCR) in a Perkin Elmer Geneamp 2400 thermal cycler with 16S rDNA (46f and 519r) primers and methyl coenzyme-M reductase (mcrAf and mcrAr) primers, respectively. 16S rDNA PCR conditions (30 cycles) were denaturing at 94° C. (30 s), annealing at 56° C. (30 s), and extension at 72° C. (1 min).

Microbial members of the consortium were characterized by cloning and sequencing the bacterial 16S rDNA and mcrA amplicons. Amplicons were purified using the Wizard PCR purification kit (Promega, Madison, Wis.) and cloned using the TA cloning kit or the Topo TA cloning kit for sequencing according to manufacturer's instructions (Invitrogen, San Diego, Calif.). Colonies were picked and 16S rRNA and mcrA gene clone fragments (133 and 48, respectively) were recovered using vector primers and mcrA primers, respectively, using PCR. For the bacterial 16S rDNA characterization, the PCR products were reamplified using 46f and 519r primers. All PCR amplicons were digested with restriction enzymes (6 μl of PCR product with 2.5 U each of MspI and HinPI) according to manufacturer's instructions (Promega, Madison, Wis.). Restriction fragments were analyzed by size separation on a 3.5% Metaphor (Cambrex, Rockland, Me.) agarose gel, restriction fragment length polymorphism (RFLP) patterns were distinguished, and the frequency with which each pattern occurred was determined.

It should be noted that the frequency of clones in the library may not correspond directly to relative phylotype numbers in the culture due to undefined differences in the number of 16S rDNA copies per cell. In addition, PCR and nucleic extraction biases may contribute to apparent differences in the abundance of RFLP patterns. Representative clones for each pattern were selected for sequencing. Amplicons to be sequenced were purified with the wizard PCR purification system, and cycle sequencing was performed on both strands using Big Dye v3.1 (Applied Biosystems, Foster City, Calif.) and run on an ABI310 genetic analyzer. Sequences were edited and assembled using Autoassembler (Applied Biosystems, Foster City, Calif.). Closest phylogenetic relatives were

determined by BLAST search of the National Center of Bioinformatics (NCBI) database (<http://www.ncbi.nlm.nih.gov/>).

Terminal restriction fragment length polymorphism (TRFLP)-PCR was performed as described above, but using 46f primer with FAM label attached. A restriction digest of 6 μl of PCR product was performed using 5 U MnlI (New England Biolabs, Beverly, Mass.). Digested samples were precipitated with 0.1 volume of 3 M sodium acetate and 2 volumes of cold 100% ethanol and resuspended in 10 μl sterile water. A 2.5 μl aliquot of the digested sample was added to 12 μl of deionized formamide and 0.5 μl ROX500 standard (Applied Biosystems). Samples were denatured at 95° C. for 5 min. DNA fragments were separated using an ABI310 sequencer (Applied Biosystem). Terminal restriction fragments were detected using 310Genescan analytical software, version 2.1.1, resulting in a TRFLP profile for each sample.

VII. Detection and Quantification of Specific Members by Quantitative PCR

Primers were used to detect organisms with abundances too low to be detected in the 16S clone library, including two known dechlorinators, *Dehalococcoides* spp., and *Desulfuromonas* spp., and methanogens. DNA copy number in an extract of WBC-2 DNA was determined by quantitative PCR (qPCR) using the quantitect SYBR green real-time PCR kit (Qiagen, Chatsworth, Calif.) and the Opticon real-time PCR system (MJ Research, now BioRad, Hercules, Calif.).

The 16S rDNA based primers used to target *Dehalococcoides* were dhc730f, 5'-GCG GTT TTC TAG GTT GTC-3' (SEQ ID NO 21) and dhc1350r, 5'-CAC CTT GCT GAT ATG CCG-3' (SEQ ID NO 22). The *Desulfuromonas* primers (designed for specificity to *Desulfuromonas* sp. strain BB1 and *D. chloroethenica* 16S rDNA) and conditions are known (Löffler et al., 2000). Methanogens were quantified using mcrA primers (Luton et al., 2002). A standard curve was determined using Ct values of serial dilutions of plasmid containing the dhc or mcrA amplified fragment, or the *Desulfuromonas* sp. strain BB1 amplicon of known concentration (and thus copy number), and the samples were plotted against that curve to determine abundance. Calculations of cell numbers were based on one 16S rDNA copy per cell for *Dehalococcoides* (www.tigr.org), and 1 mcrA copy per cell for methanogens. For the purposes of calculating cell numbers, nucleic acid extractions were assumed to be perfect, because no measurement of extraction efficiency is available. For microscopic counts, culture samples were suspended in 0.01% Triton X-100 and stained with 5 $\mu\text{g}/\text{ml}$ 4',6-diamidino-2-phenylindole (DAPI) in 1 \times phosphate-buffered saline (PBS), filtered onto a black Nucleopore filter (0.2 μm), and viewed using epi-fluorescence.

Although the present invention has been described in terms of particular exemplary and alternative embodiments, it is not limited to those embodiments. Alternative embodiments, examples, and modifications which would still be encompassed by the invention may be made by those skilled in the art, particularly in light of the foregoing teachings.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 22

<210> SEQ ID NO 1

<211> LENGTH: 425

<212> TYPE: DNA

<213> ORGANISM: Unknown

<220> FEATURE:

-continued

<223> OTHER INFORMATION: uncultured bacterium

<400> SEQUENCE: 1

```

gcgagatgat ctcttcgggg agattctagc gccggacggg tgagtaacac gtgggtaacc    60
tgoctcaaag aggggatag cctcccgaaa gggagattaa taccgataa cattattttc    120
tcgcatgaga agataatcaa aggagcaatc cgctttgaga tggaccocgc gcgcattagc    180
tagttggtga ggtaacggct caccaaggcg acgatgcgta gccgacctga gagggtgatc    240
ggccacattg gaaactgagc acggtccaga ctectacggg aggcagcagt ggggaatatt    300
gcgcaatggg ggaaccctg acgcaagcaa cgcgcgctga gtgatgaagg tcttcggatt    360
gtaaagctct gtctttgggg acgataatga cggtaaccac ggaggaagcc accgctaact    420
acgtg                                           425

```

<210> SEQ ID NO 2

<211> LENGTH: 430

<212> TYPE: DNA

<213> ORGANISM: Unknown

<220> FEATURE:

<223> OTHER INFORMATION: uncultured bacterium

<400> SEQUENCE: 2

```

agtcgagcga gatgatccct ccgggagatt ctaccggcgg acgggtgagt aacgcgtggg    60
taacctgcc tatgaaagg aatagcctcg gaaaactggg agtaatgcct tataatacat    120
tttggtcgca cgaactatgat gttaaacgct ccggtgccat aggatggacc cgcgtcccat    180
tagctagtgt gtgagataac agcccaccaa ggogaogatg ggtaacgggg tctgagaggg    240
cgaacggcca cactggaact gagacacggt ccagactcct acgggaggca gcagtgggga    300
atattgcgca atgggggcaa ccctgacgca gcaataccgc gtgagtgaag aaggttttcg    360
gatcgtaaag ctctgttatt ggggaagaaa aaagacggta cccaagaaga aagtccgggc    420
taactacgtg                                           430

```

<210> SEQ ID NO 3

<211> LENGTH: 478

<212> TYPE: DNA

<213> ORGANISM: Unknown

<220> FEATURE:

<223> OTHER INFORMATION: uncultured bacterium

<400> SEQUENCE: 3

```

agtcgaacgg agtaataaaa taagtttact tagattatta cttagtggcg aacgggtgag    60
taacgcgtgg gtaacctgcc cttaaagacc ggacaacagc tggaaacggc tgctaatacc    120
ggatggattt attggaagcg atcttcta ataggaaagct ggcctctgta tatgctagcg    180
cttagggatg gatccgcgto tgattagota gttggtaggg taatggccta ccaaggcgac    240
gatcagtagc cggcctgaga gggtaaaccg ccacactggg actgagacac ggcccagact    300
cctacgggag gcagcagtg ggaatcttc gcaatggagc aaagtctgac ggagcaacgc    360
cgcgtgtatg aagaaggcct tcgggttgta aaactactgt gttggggaag aacggctgga    420
gtgtaataaa tgcttcagat tgaocgtaac caacgaggaa gccccggota actacgtg    478

```

<210> SEQ ID NO 4

<211> LENGTH: 447

<212> TYPE: DNA

<213> ORGANISM: Unknown

<220> FEATURE:

<223> OTHER INFORMATION: uncultured bacterium

-continued

<400> SEQUENCE: 4

```

ggggcagcac aaggagcaat ctgaggtggc gaccggcgca cgggtgagta acacgtatgc   60
aacctgtctg taagcgggga ataaccggtt gaaagacgga ctaataccgc ataatactgg   120
agatctgoat ggatattcag ttaaaccattt atggottaca gatgggcatg cgcattgatta   180
gatagttgga gagtaaccg ccccccaagt caacgatcat taggggttct gagaggaagg   240
tccccacac tggctactgag acacggacca gactcctacg ggaggcagca gtgaggagta   300
ttggtcaatg ggcgagagcc tgaaccagcc aagtcgcgtg caggaagaat gtctatgga   360
ttgtaaactg cttttgcagg ggaataaagt gagccaccgt gtggtttttt gtatgtaote   420
tgcgaataag gatcggctaa ctccgtg                                     447

```

<210> SEQ ID NO 5

<211> LENGTH: 454

<212> TYPE: DNA

<213> ORGANISM: Unknown

<220> FEATURE:

<223> OTHER INFORMATION: uncultured bacterium

<400> SEQUENCE: 5

```

ggggcagcac gggagtagca atacaactgg tggcgaccgg cgcacgogtg agtaaccgct   60
atgcaaccta cctatcagag gggataaacc cggcgaaagt cggactaata ccgcataaaa   120
cagggggcgc cgcattggtg tatttgttaa agaattcgtc gatagatggg catgcgttcc   180
attagggtag ttggtgaggt aacggctcac caagccgacg atggataggg gaactgagag   240
gttggtcccc cacactggta ctgagacacg gaccagactc ctacgggagg cagcagtgag   300
gaatattggt caatggggca gagcctgaac cagcccaagt cgcgtgaagg aagaaggatc   360
tatggtttgc taaactctt tttgcagggg aataaagtgc gggacgtgto ctgtttcgta   420
tgtaccctga gaataaggat cggctaactc cgtg                                     454

```

<210> SEQ ID NO 6

<211> LENGTH: 464

<212> TYPE: DNA

<213> ORGANISM: Unknown

<220> FEATURE:

<223> OTHER INFORMATION: uncultured bacterium

<400> SEQUENCE: 6

```

acggagtgaa ggagcttget ctttcattta gtggcgcacg ggtgagtaac gcgtagataa   60
tctgccttag actctgggat aacatctcga aaggggtgct aataccggat aagcccacga   120
tggcgtaagt cattcgggga aaagggggcc tctgaatatg ctcttgatct aagatgagtc   180
tgcgtaccat tagctagttg gtagggtaag agcctaccaa ggcgacgatg gttagctggt   240
ctgagaggat gatcagccac actggaactg agacacggtc cagactocta cgggaggcag   300
cagtggggaa ttttgcgcaa tgggggaaac cctgacgcag caacgcgcgg tyagtgatga   360
aggctttcgg gtctgaaagc tctgtctaga gggaagaat gataatcggg taatacccgg   420
ttttcttgac ggtacctctg aaggaagcac cggctaactc cgtg                                     464

```

<210> SEQ ID NO 7

<211> LENGTH: 446

<212> TYPE: DNA

<213> ORGANISM: Unknown

<220> FEATURE:

<223> OTHER INFORMATION: uncultured bacterium

<400> SEQUENCE: 7

-continued

```

gcgatgaag ggagcttget ctctgattoa gggggggaag ggtgagtaat gcttaggaat 60
ctgcctgata gtgggggaca acgttttgaa aggaacgcta ataccgata cgtctacgg 120
gagaaagcag gggaccttgg ggccttgcgc taccagatga gcctaggteg gattagctag 180
ttggtgaggt aacggctcac caaggcgacg atccgtaact ggtctgagag gatgatcagt 240
cacactgaa ctgagacacg gtccagactc ctacgggagg cagcagtggg gaattatgga 300
caatgggoga aagcctgac cagccatgcc gogtgtgtga agaaggtctt cggattgtaa 360
agcaacttaa gttgggagga agggcattaa cctaatacgt tagtgltttg acgttacga 420
cagaataagc accggctaac ttcgtg 446

```

```

<210> SEQ ID NO 8
<211> LENGTH: 433
<212> TYPE: DNA
<213> ORGANISM: Unknown
<220> FEATURE:
<223> OTHER INFORMATION: uncultured archaeon

```

<400> SEQUENCE: 8

```

tgcatacacc gatgacatcc tcgacaacaa cgtgtactac gacgttgact acatcaacga 60
caagtacaac ggtgcagcaa ccacggcaa ggacaacaag atcaaggcca cctcgcagct 120
cgtaaaggac atcgcaaccg agtccactat ctaccgtatc gagacctacg agaagttccc 180
gactgocctt gaagaccact tcggtggatc ccagagagca accgtgctcg cagctgcagc 240
cgggtgcgca acctccctcg caaccgcaa tcgaaatgct ggtctctccg gctggtacct 300
ctccatgtac ctgcacaagg aagcatgggg ccgtctcggc ttctctggat acgacctgca 360
ggaccagtgc ggtgcacaaa acgtctctgc ctaccagggc gacgaaggtc tcccagagca 420
actccgtggt cca 433

```

```

<210> SEQ ID NO 9
<211> LENGTH: 415
<212> TYPE: DNA
<213> ORGANISM: Unknown
<220> FEATURE:
<223> OTHER INFORMATION: uncultured archaeon

```

<400> SEQUENCE: 9

```

agcatcaccc gacaacatcc tcgacaggtt cacctactac ggtatggact acctaaagga 60
caagtaecgc ggctactcgc aggcaccagc aaccacggag gtctgcaacg acctcgcaac 120
tgaggtcacg ctgaacgcca tggagcagta ccagcagttc ccaaccatga tggaggacca 180
cttcggcggg tcccagcgtg ctggtgtcat tgcagcagca tcaggtttga ccaactgcaat 240
cggtaaccgg aactccaatg ccggtctgaa ccgctggtac ctctcgatgc tcatgcacaa 300
ggaaggatgg tcacgtctcg gttctctcgg ctaccacctg caggaccagt gtggttcagc 360
aaactcgatg tcgatcagac cggatgaagg tctctcggg gaactccgtg gacca 415

```

```

<210> SEQ ID NO 10
<211> LENGTH: 432
<212> TYPE: DNA
<213> ORGANISM: Unknown
<220> FEATURE:
<223> OTHER INFORMATION: uncultured archaeon

```

<400> SEQUENCE: 10

```

tgcatacacc gatgacatcc tcgacaacaa cgtgtactac gacgttgact acatcaacga 60
caagtacaac ggtgctgcaa ccacggcaa ggacaacaaa gtaaaggcaa cctcgcagct 120

```

-continued

```

cgtaaaggac atcgcaaccg agtccactat ctacgggtatc gagacctacg agaagttccc 180
gactgccctt gaagaccact tcggtggatc ccagagagca accgtgctcg cagctgcagc 240
cgggtgcgca acctccctcg caaccgcaaa cgccaacgct ggtctctcgg gctggtacct 300
ctccatgtac ctgcacaagg aagcatgggg cgtctcggct tcttcggata cgacctgcag 360
gaccagtgcg gtgccacaaa cgttctgtcc taccagggcg acgaaggtct cccagacgaa 420
ctccgtggtc ca 432

```

```

<210> SEQ ID NO 11
<211> LENGTH: 432
<212> TYPE: DNA
<213> ORGANISM: Unknown
<220> FEATURE:
<223> OTHER INFORMATION: uncultured archaeon

```

```

<400> SEQUENCE: 11

```

```

tgcatacacc gatgacatcc tcgacaacaa cgtgtactac gacgttgact acatcaacga 60
caagtacaac ggtgctgcaa ccaccggcaa ggacaacaaa gtaaaggcaa cctcgcagct 120
cgtaaaggac atcgcaaccg agtccactat ctacgggtatc gagacctacg agaagttccc 180
gactgccctt gaagaccact tcggtggatc ccagagagca accgtgctcg cagctgcagc 240
cgggtgcgca acctccctcg caaccgcaaa cgccaacgct ggtctctcgg gctggtacct 300
ctccatgtac ctgcacaagg aagcatgggg cgtctcggct tcttcggata cgacctgcag 360
gaccagtgcg gtgccacaaa cgttctgtcc taccagggcg acgaaggtct cccagacgga 420
ctccgtggtc ca 432

```

```

<210> SEQ ID NO 12
<211> LENGTH: 433
<212> TYPE: DNA
<213> ORGANISM: Unknown
<220> FEATURE:
<223> OTHER INFORMATION: uncultured archaeon

```

```

<400> SEQUENCE: 12

```

```

tgcatacact gatgacatcc tcgacaadaa cgtgtactac gacgttgact acatcaacga 60
caagtacaac ggtgctgcaa cagtcggcaa ggacaacaag gtaaaggcaa cctcgaagt 120
cgtaaaggac atcgcaaccg agtccactat ctacgggtatc gagacctacg agaaattccc 180
aacgcctt gaagaccact tcggtggatc ccagagagca accgtgctcg cagctgcagc 240
cgggtgcgca tgtgcactcg gaactgcaaa cgccaatgct ggtctctcag gctggtacct 300
ctccatgtac ctgcacaagg aagcatgggg cagactcggc ttcttcggat acgacctgca 360
ggaccagtgc ggtgcacaaa acgttctgtc ctaccagggc gacgaaggtc tcccagacga 420
actccgtggt cca 433

```

```

<210> SEQ ID NO 13
<211> LENGTH: 415
<212> TYPE: DNA
<213> ORGANISM: Unknown
<220> FEATURE:
<223> OTHER INFORMATION: uncultured archaeon

```

```

<400> SEQUENCE: 13

```

```

tgcatacacc gacaacatcc tcgatgacta caccaggtac ggtattgact acgtcaagaa 60
gaaacacggg ggcacggca aggcacagtc aaccacaggaa gttgtcagcg acattgcaac 120
cgaggtcaac ctctacggta tggagcagta cgaaacagtc ccgaccgcc tcgagagcca 180

```

-continued

```

cttcggtgga tcccagcgtg cgtctgtcct tgcagcagca tcaggtatct cctgttcaat 240
ggcaacggca aactccaacg ctggcctgaa cggctggtag ttgtccatgc tcatgcacaa 300
ggaaggctgg tcacgtcttg gcttctctgg ctacgacctg caggaccagt gcggttcgcg 360
aaactccatg tcgatccgtc ccgacgaggg attactcggc gagctccgtg gaccg 415

```

```

<210> SEQ ID NO 14
<211> LENGTH: 437
<212> TYPE: DNA
<213> ORGANISM: Unknown
<220> FEATURE:
<223> OTHER INFORMATION: uncultured archaeon

```

```

<400> SEQUENCE: 14

```

```

tgcatacacc gacaacatcc tcgatgacta cacctactac ggtatggatt acatcaagga 60
caagtacaaa gtcaactgga agagcccgtc agccaaggac aaggtaagc caaccagga 120
tctcgtcaat gagctcgeta ccgaagtac actctacggt atggagcagt acgagcagtt 180
cccgaccctt attggaagacc acttcggtag atcccagcgt gctggtgttc tcgagctgc 240
atccggtctt acctgtgcaa tcgccaccgg taactccaac gctggcctga acggctggta 300
cctgtccatg cttgcccaaa aggaaggctg gtcacgtctc ggcttctctg gctacgacct 360
tcaggaccag tgcggttcgg caaactcgtc ctctatcaga cctgacgaag gctgtatcgg 420
cgagctccgt ggaaccg 437

```

```

<210> SEQ ID NO 15
<211> LENGTH: 415
<212> TYPE: DNA
<213> ORGANISM: Unknown
<220> FEATURE:
<223> OTHER INFORMATION: uncultured archaeon

```

```

<400> SEQUENCE: 15

```

```

agcatacacc gacaacatcc tcgacgagtt cacctactac ggtatggact acttaaagga 60
caagtaaggc ggctactcgc aggcaccagc aaccaggag gtctgtaac acctcgaac 120
tgaggtcagc ctgaacgcaa tggagcagta cagcagctc ccaaccatga tggaggacca 180
cttcggcggt tcccagcgtg ctggtgtcat tgcagcagca tcaggtttga ccactgcaat 240
cggtaccggg aactccaatg ccggtctgaa cggctggtag ctctcagatgc tcatgcacaa 300
ggaaggctgg tcacgtctcg gttctctcgg ctacgacctg caggaccagt gtggttcagc 360
aaactcagat tcgatccagc cggatgaagg tctctctcgg gaaactccgtg gacca 415

```

```

<210> SEQ ID NO 16
<211> LENGTH: 415
<212> TYPE: DNA
<213> ORGANISM: Unknown
<220> FEATURE:
<223> OTHER INFORMATION: uncultured archaeon

```

```

<400> SEQUENCE: 16

```

```

tgcatacacc gacaacatcc ttgatgacta caccagtagc ggtctggact atatcaagaa 60
gaaacacggt ggctcagcaa aggcacaggc gaccaggaa gtctcagcg acatcgaac 120
cgaggtcaac ctctacgcta tggagcagta cgaacacctc ccaaccgctc tcgagagcaa 180
cttcggtgga tcccagcgtg catccgtcct tgcagcagca tcgggtatca ccacttcaat 240
ggctaccgca aactccaacg cgggcctgaa cggctggtag ttgtccatgc tcatgcacaa 300
ggaaggctgg tcacgtctcg gttctctcgg ctacgacctg caggaccagt gtggttcgcg 360

```

-continued

 aaactccatg tcgatccgtc cggagcaggg attactcggc gagctccgtg gaccg 415

<210> SEQ ID NO 17
 <211> LENGTH: 415
 <212> TYPE: DNA
 <213> ORGANISM: Unknown
 <220> FEATURE:
 <223> OTHER INFORMATION: uncultured archaeon

<400> SEQUENCE: 17

agcctacacc aacgatgtcc tggacgactt ctgctactac ggcgtogact tcgccgcaga 60
 taagttcgga ggattcgcca agtcacccaa gactctggac atcgccaagg agctggccac 120
 tgaggtcaac gctacaggca tggagcagta cgaggaatcc cggactctgc tcgaggatca 180
 cttegggtgga tcccagagag catccgttct ggcagccgca tcgggtatca cttecccat 240
 cgcctctggc cacagccagg taggccttgc cggatggtac ctgtctatgc tcctgcacaa 300
 ggaaggctgg ggaagcctgg gctctctcgg ctacgacctg caggatcaat cgggtccaac 360
 caacgtgttc tctaccagt cggagcaggg cgcacccctc gagctgaggg gagct 415

<210> SEQ ID NO 18
 <211> LENGTH: 435
 <212> TYPE: DNA
 <213> ORGANISM: Unknown
 <220> FEATURE:
 <223> OTHER INFORMATION: uncultured archaeon

<400> SEQUENCE: 18

tgcatacacc gatgatatcc tcgacaacaa catctactac gacgttgact acatcaacga 60
 caagtacaac ggtgcagcaa ccctcggcaa ggacaacaag ttaaaggcaa gcctcgyaat 120
 cgtaaaggac attgcaaccg agtcacacat ctacgggatg gagaactacg agaagttccc 180
 aaccgctctt gaagaccact tcggtggatc ccagagagca accgtgctcg cagctgcagc 240
 tgggtgttga tgcgcctcgg gaadtgcaaa cgcacaacgt ggtctctcgg gctggtacct 300
 ctccatgtac ctgcacaagg aagcatgggg ccgtctcggg tottcggata cgacctgca 360
 ggaccagtgc ggtgcacaaa acgtttctgt cctaccaggg cgaagcaagg ctcccagagc 420
 aactccgtgg tccca 435

<210> SEQ ID NO 19
 <211> LENGTH: 367
 <212> TYPE: DNA
 <213> ORGANISM: Unknown
 <220> FEATURE:
 <223> OTHER INFORMATION: uncultured archaeon

<400> SEQUENCE: 19

tctacgtcaa gagaaacatg gtggcatcgc aaaggcaag gcgacacagg aagttgtcag 60
 cgacatcgca accgaggtca acctctacgg tatggagcag taagagacct acccgaccgc 120
 cctcgagagc cacttcggcg gatcccagcg tgcacccgct cttgcagcag catcaggtat 180
 ttctgttga atggcaaccg caaactccaa cgcgggtctg aacggctggt acctgtcgat 240
 gctcatgcac aaggaaggct ggtcacgtct cggtttcttc ggctacgacc tgcaggacca 300
 gtgcggttcc gcaactccaa tgtcaatccg tccagacgag ggattactcg gagaactccg 360
 tggaccg 367

<210> SEQ ID NO 20
 <211> LENGTH: 415
 <212> TYPE: DNA

-continued

```

<213> ORGANISM: Unknown
<220> FEATURE:
<223> OTHER INFORMATION: uncultured archaeon

<400> SEQUENCE: 20
tgcatacacc aacgatgtcc tggacgactt ctgctactac ggcgtcgact tcgcccaga    60
caagtteggg ggattcgcca aggcacccaa gacctggat atcgccaagg aattggcaac    120
cgaggccaac gcttatgggt ttagcagta cgaggcattc ccgactctgc tcgaggatca    180
cttcgggtgga tcccagaggg catccgtcct cgcagccgca tcgggtatca cctcagccat    240
cgctcccgcc caccagccagg tcggtctcgc cggctggtac ctgagcatgc tctctcacia    300
ggaatcctgg ggacgcttgg gctttctgg ctacgacttg caggatcaat cgggtccaac    360
caacgtattc tctaccagt cagacgaggg caaccocgto gagctgaggg gcgca        415

<210> SEQ ID NO 21
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 21
gcggtttttt aggttgtc        18

<210> SEQ ID NO 22
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 22
caccttgctg atagggg        18
    
```

What is claimed is:

1. A method for dechlorinating a mixture of chlorinated ethanes and chlorinated ethenes, comprising:
 - contacting a mixture of chlorinated ethanes and chlorinated ethenes with a microbial composition comprising an isolated bioremediative consortium comprising strains of microorganism comprising *Clostridium*, *Acetobacter*, *Dehalobacter*, *Bacteroides*, and *Proteobacteria*; and
 - concurrently anaerobically dechlorinating the mixture of chlorinated ethanes and chlorinated ethenes.
2. A method for dechlorinating a mixture of chlorinated ethanes and chlorinated ethenes, comprising:
 - contacting a mixture of chlorinated ethanes and chlorinated ethenes with a microbial composition comprising an isolated bioremediative consortium comprising strains of microorganism comprising *Clostridium*, *Acetobacter*, *Dehalobacter*, *Bacteroides*, *Proteobacteria*, and *Methanomicrobia*; and
 - concurrently anaerobically dechlorinating the mixture of chlorinated ethanes and chlorinated ethenes.

3. A method according to claim 1, wherein the chlorinated ethanes comprise at least one of 1,1,2-tetrachloroethane; 1,1,2-trichloroethane; 1,2-dichloroethane, or chloroethane.
4. A method according to claim 1, wherein the chlorinated ethenes comprise at least one of cis 1,2-dichloroethene; trans 1,2-dichloroethene; vinyl chloride; or tetrachloroethene.
5. A method according to claim 1, wherein the mixture further comprises chlorinated methane.
6. A method according to claim 5, wherein the chlorinated methane comprises carbon tetrachloride or chloroform.
7. A method according to claim 1, wherein the microbial composition further comprises *Dehalococcoides* and at least one of *Methanosarcina* or *Methanosaeta*.
8. A method according to claim 1, wherein the microbial composition comprises strains of microorganism comprising *Clostridiales*, *Cytophaga-flavobacterium-bacterioides*, *Proteobacteria*, and *Methanomicrobia*.
9. A method according to claim 1, wherein the chlorinated ethane comprises 1,1,2,2-tetrachloroethane and said anaerobically dechlorinating occurs in the presence of tetrachloroethene.

* * * * *